

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 1/00, C07H 21/02	A1	(11) International Publication Number: WO 99/50280 (43) International Publication Date: 7 October 1999 (07.10.99)
(21) International Application Number: PCT/US99/07150 (22) International Filing Date: 31 March 1999 (31.03.99) (30) Priority Data: 60/080,146 31 March 1998 (31.03.98) US (71) Applicant (for all designated States except US): TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA [US/US]; Suite 300, 3700 Market Street, Philadelphia, PA 19104-3147 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): EL-DEIRY, Wafik, S. [US/US]; Apartment P113, 1500 Locust Street, Philadelphia, PA 19102 (US). WEBER, Barbara, L. [US/US]; 331 Mallwyd Road, Merion, PA 19066 (US). (74) Agents: REED, Janet, E. et al.; Dann Dorfman Herrell and Skillman, Suite 720, 1601 Market Street, Philadelphia, PA 19103 (US).		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: COMPOSITIONS AND METHODS FOR CONTROLLING BRCA1-MEDIATED P53-DEPENDENT AND -INDEPENDENT REGULATION OF TRANSCRIPTION (57) Abstract It has been discovered that BRCA1 mediates transcriptional activation of <i>p21</i> and other tumor suppressor genes in a p53-dependent and p53-independent manner. Assays and compositions for identifying compounds that enhance or repress cellular proliferation via these newly-discovered BRCA1-mediated pathways are disclosed. Diagnostic methods and methods of controlling cell proliferation by altering the pathways are also disclosed.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**COMPOSITIONS AND METHODS FOR CONTROLLING
BRCA1-MEDIATED P53-DEPENDENT AND
-INDEPENDENT REGULATION OF TRANSCRIPTION**

This application claims priority to U.S. Provisional Application Serial No. 60/080,146, filed March 31, 1998, incorporated by reference herein.

5 Pursuant to 35 U.S.C. §202(c), it is acknowledged that the U.S. Government has certain rights in the invention described herein, which was made in part with funds from the National Institutes of Health, Grant No. CA57601.

10

FIELD OF THE INVENTION

This invention relates to regulation of cell proliferation. In particular, this invention provides plasmids, cell lines, kits and assay methods for
15 identifying compounds that affect p53-dependent and p53-independent regulation of transcription by BRCA1, as well as diagnostic and therapeutic methods relating thereto.

BACKGROUND OF THE INVENTION

20 Various scientific and scholarly articles are referred to in parentheses throughout the specification. These articles are incorporated by reference to describe the state of the art to which this invention pertains.

Breast cancer is one of the most common
25 malignancies affecting women. About 5-10% of all cases are estimated to be familial. Susceptibility To
~~early-onset breast and ovarian cancer is conferred by~~
mutations in a gene on chromosome 17q21. This gene
termed *BRCA1*, has been identified by positional cloning
30 (Miki et al., 1994, Science 266:66-71). Mutations in the
BRCA1 gene account for about 50% of inherited breast

cancer cases and 80% of families predisposed to both breast and ovarian cancer.

Much effort has been recently directed at understanding the biochemical function of *BRCA1*. The human *BRCA1* gene encodes an 1863 amino acid nuclear protein that is expressed in a wide variety of adult human tissues (Marquis et al., 1995, Nat. Genet. 11:17-26). Evidence implicates a role for *BRCA1* in the control of gene expression. *BRCA1* contains a nuclear localization signal (Thakur et al., 1997, Mol. Cell. Biol. 17:444-452), a C-terminal domain that transactivates gene expression when fused to a heterologous DNA binding domain (Chapman and Verma, 1996, Nature 382:678-679; Monteiro et al., 1996, PNAS 93:13595-13599) and *BRCA1* has been found as a component of RNA polymerase II (Scully et al., 1997, PNAS 94:5606-5610).

Several lines of evidence suggest that *BRCA1* protein may play an important role as a suppressor of cell proliferation. Antisense oligonucleotides to *BRCA1* mRNA accelerated the growth of normal and malignant mammary epithelial cells (Thompson et al., 1996, Nat. Genet. 9:444-450). Similarly, NIH 3T3 cells with a stable expression of antisense *BRCA1* mRNA showed accelerated growth rate, anchorage independent growth, and tumorigenicity in nude mice unlike the parental cell line and stable cell lines expressing sense *BRCA1* mRNA (Rao et al., 1996, Oncogene 12:523-537). In addition, transfection of the wild-type *BRCA1* gene inhibited the growth of breast and ovarian cancer cell lines (Holt et al., 1996, Nat. Genet. 12:298-302). Retroviral introduction of wild-type *BRCA1* gene into the already established MCF7 human breast tumors inhibited the growth of the tumors significantly and also increased the

survival of mice (Holt et al., 1996, *supra*). Another evidence that *BRCA1* may have an antiproliferative function is that human breast cancers involving mutations in *BRCA1* feature increased proliferation, aneuploidy, 5 high S-phase fraction and high mitotic grade (Marcus et al., 1997, *Cancer* 80:543-556).

Evidence also suggests that *BRCA1* may activate cell proliferation. The protein p21 is a universal cell-cycle inhibitor that specifically binds cyclin-CDK 10 complexes and proliferating cell nuclear antigen, thereby serving as a potent inhibitor and effector of cell-cycle checkpoints (Sherr and Roberts, 1995, *Genes Dev.* 9:1149-1163). Cells from *BRCA1*-null mouse embryos have increased levels of p21 mRNA, which suggests that *BRCA1* 15 may suppress p21 expression during development to allow cell growth (Hakem et al., 1996, *Cell* 85:1009-1023).

A number of observations have also linked *BRCA1* to DNA damage response pathways (Brugarolas and Jacks, 1997, *Nature Med.* 7:721-722). *BRCA1* normally co- 20 localizes with Rad51, the human homolog of the *Escherichia coli* RecA protein, at nuclear dot structures that may be sites of checkpoint processing in S-phase cells (Scully et al., 1997, *Cell* 88:265-275; Tashiro et al., 1996, *Oncogene* 12:2165-2170). Following exposure to 25 DNA damaging agents, *BRCA1* becomes hyperphosphorylated and disperses from dot structures and then dynamically accumulates at PCNA-containing replication structures, suggesting a role in the checkpoint response (Scully et al., 1997, *Cell* 90:425-435).

30 It is not known how *BRCA1* affects gene transcription in normal cells and how mutations in *BRCA1* cause cells to become cancerous. Lack of this information hinders the development of methods of

treatment of mutant BRCA1-induced tumors. In particular, it is very desirable to have a model system of BRCA1 activation of gene transcription so that alternate ways of activating the tumor-suppressing genes can be devised
5 for tumor cells.

SUMMARY OF THE INVENTION

In accordance with it the present invention, two ways in which BRCA1 inhibits cell proliferation have
10 been discovered. First, BRCA1 inhibits S-phase cell cycle progression by transactivating expression of the CDK-inhibitor p21^{WAF1/CIP1} in a p53-independent manner. Second, BRCA1 interacts with p53 *in vitro* and *in vivo*, and increases p53-dependent transcription from the p21
15 and bax promoters. These pathways of BRCA1 action provide novel targets to which pharmaceutical agents capable of affecting cell proliferation and apoptosis can be isolated, and can then be used in the treatment and control of cellular proliferation disorders.

20 One aspect of the present invention is drawn to compositions and methods for use in assay systems to identify such novel pharmaceutical agents. One method uses an *in vitro* or cell-based system that measures the ability of a test compound to enhance or interfere with
25 the association between BRCA1 and p53. This method comprises allowing BRCA1 and p53 to associate in the presence or absence of the test agent, separating the BRCA1/p53 complexes, and determining the amount of complex formed in the presence, as compared to the
30 absence, of the test compound.

Another method determines the effect of the pharmacological agent on p53-dependent and -independent BRCA1-mediated gene transcription. This method comprises

the culturing of various cell lines containing reporter gene(s) driven by the *p21* or *bax* promoters with and without the agent and assaying the activity of the reporter genes. Using appropriate cell types,
5 e.g., BRCA1-/BRCA1- and WT cell lines, this method can be used to determine the ability of a pharmacological agent to coactivate p53 in a manner similar to BRCA1 coactivation of p53.

A variation of the above-described method
10 determines the effect of a pharmacological agent on p53-dependent and p53-independent BRCA1 activated apoptosis or cell cycle arrest. In a preferred embodiment, this method comprises the culturing of p53⁺/p53⁺ and p53⁻/p53⁻ cell lines with and without the agent and assaying for
15 apoptosis or cell cycle arrest, using standard methods.

According to another aspect of the present invention, isolated nucleic acids and/or polypeptides are provided, corresponding to (1) the BRCA1/p53 protein binding domains; and (2) the *p21* transcriptional
20 regulatory region required for BRCA1-mediated, p53-independent transactivation, which have been identified in accordance with the present invention. In a preferred embodiment, the nucleic acid molecule encodes, or the polypeptide comprises, residues 224-500 of the BRCA1
25 protein. In another preferred embodiment, the nucleic acid molecule encodes, or the polypeptide comprises, residues 300-395 of the p53 polypeptide. In another preferred embodiment, the portion of the *p21* promoter comprising the transcription start site and 5' sequence
30 to -143, specifically the region 93 to 143 nucleotides upstream of the transcription start site, is provided.

The aforementioned isolated nucleic acids or peptides are useful for some of the assays described

above, as well as for diagnostic assays to determine the predisposition of an individual to developing certain tumors, based on the presence or absence of functional mutations that disrupt BRCA1-mediated control of cell proliferation through p21, bax or other downstream targets, either dependent on or independent of p53.

According to other aspects of the invention, methods are provided to control cell proliferation, by up-regulating or down-regulating p53 dependent or independent BRCA1-mediated transcriptional activation of p21 or other downstream targets involved in apoptosis or cell cycle arrest. Up-regulation of these pathways should result in suppression of cell proliferation either by inducing cell cycle arrest or apoptosis, thereby providing a useful cancer therapy. Down-regulation of these pathways should result in increased cell proliferation, which may be useful in cell culture for expansion of recalcitrant cell lines, or which may be useful as a research tool.

In another aspect of the present invention, kits are provided for practicing the assays and diagnostic methods set forth above. These kits contain one or more plasmids, cell lines, antibodies, reagents and the like, along with instructions for performing the assays or diagnostic methods.

Other features and advantages of the present invention will be better understood by reference to the drawings, detailed description and examples that follow.

30 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. BRCA1 transfection inhibits DNA synthesis in human cancer cells. SW480 cells were cotransfected with pCR3, pCR3-BRCA1, pCEP4, or pCEP4-p53.

The percent BrdU(+)/GFP(+) cells shown was determined by analyzing at least 75 GFP(+) cells for each transfection in three fields, for each of two independent experiments.

Figure 2. BRCA1 transactivates the human and mouse p21-promoter and upregulates endogenous p21 protein expression. **Fig. 2A.** SW480 or HCT116 cells were cotransfected with pWWP-Luc and either pCR3 or pCR3-BRCA1 and luciferase activity was measured 24 hrs later as described in Example 1. **Fig. 2B.** CV1 cells were cotransfected pCAT1 and pCR3 or pCR3-BRCA1 and CAT activity was measured after 48 hrs later as described in Example 1. **Fig. 2C.** Structure of the human p21-promoter luciferase reporter and several 5'-deletions are shown schematically (left). The 5' end of each deletion is as indicated while the 3' boundary is 16 bp downstream of the p21 transcription initiation site (El-Deiry et al., 1993, Cell 75:817-825), fused to the luciferase reporter gene (El-Deiry et al., 1995, Cancer Res. 55:2910-2919). S1 and S2 indicate the relative locations of the two p53 DNA-binding sites within the 2.3 kb regulatory region upstream of the *WAF1/CIP1* gene (El-Deiry et al., 1995, *supra*). pWWP-Luc or the 5'-deletion mutants were cotransfected with pCR3 or pCR3-BRCA1 into SW480 cells and luciferase activity was assayed as in Fig. 2A.

Figure 3. BRCA1 mutants are defective for activation of p21. **Fig. 3A.** Structure of BRCA1 and various mutants used is shown schematically (left). Synthetic mutants lacking the functional nuclear localization signal (Δ NLS) (El-Deiry et al., 1995, *supra*), a Δ NLS mutant with a C-terminal fused NLS (Δ NLS/C+NLS), mutants lacking the RAD51-interacting domain (Δ 515-1091), the C-terminal transactivation domain (Δ 1314-1863) or both (Δ 500-1863) are shown. The

corresponding extent of p21 induction (data from Figs. 3B and 3C) is shown (right; "++" = greater than 5-fold induction; a single arrow = 1.5- to 2-fold reduction in p21 activation; double arrow = substantial decrease in p21 activation, corresponding to less than 1.5-fold induction). Fig. 3B, COS-7; or Fig. 3C, SW480 cells were co-transfected with pWWP-Luc and pCR3 or either WT or mutant BRCA1 expression plasmids as indicated and luciferase activity was determined as in Fig. 2.

10 **Figure 4.** BRCA1 fails to inhibit DNA synthesis in p21^{-/-} human cancer cells. The percent BrdU(+)/GFP(+) cells was determined following transfection of p21^{+/+} (lanes 1,2) or p21^{-/-} (lanes 3,4) HCT116 cells with pCR3 (lanes 1,3) or BRCA1 (lanes 2,4) as described in Fig.1.

15 **Figure 5.** BRCA1 stimulates p53-dependent transcription. Fig. 5A, SW480; Fig. 5B, MEF p53^{-/-}; or Fig. 5C, HCT116 cells were cotransfected with different combinations (as indicated by a A+") of BRCA1 or its vector, pCR3 (3.5 µg) and p53 or its vector, pCEP4. (2 µg) and PG13-LUC reporter (1 µg). Fig. 5D. MCF7 cells were transfected by either BRCA1 or its vector, pCR3 (16 µg) and PG13-LUC reporter (4 µg). Fig. 5E. SW480 cells were transfected with the PG13-LUC reporter (0.5 µg) and increasing concentrations of BRCA1 or its vector, pCR3 (1, 2, and 3.5 µg, respectively) either in the presence or absence of p53 (0.1 µg). Fig. 5F and Fig. 5G. SW480 cells were cotransfected with different combinations (as indicated "+ ") of BRCA1 or its vector, pCR3 (3.5 µg) and p53 or its vector, pCEP4 (2 µg) along with 1 µg of either
20 the p21-promoter WWP-LUC reporter (Fig. 5F) or the bax-promoter pbax-LUC reporter (Fig. 5G). Cells were harvested after 24 hrs of transfection and luciferase activity was measured as described in Example 2.

Figure 6. Schematic diagram identifying the p53-binding region of human BRCA1. The structure of BRCA1 protein is shown, including the nuclear localization signal (NLS), the transactivation domain (TA), the RING-finger domain (RING), the Rad51-interacting domain (Rad51) and the p53 binding domain (p53 binding). Mutants of BRCA1 tested for interaction with p53 are shown along with their ability to interact with p53 protein *in vitro* ("+" indicates interaction while "-" indicates no interaction).

Figure 7. Schematic diagram of human p53 identifying the BRCA1-interacting domain. The structure of the p53 protein is shown including the transactivation domain (TA), the DNA-binding domain, the tetramerization domain (TD) and the BRCA1-binding domain (bracket). Truncation mutants of p53 are shown along with their ability to interact with BRCA1 *in vitro* (right; "+" indicates interaction while "-" indicates no interaction).

Figure 8. Transactivation-deficient BRCA1 mutants are defective in stimulation of p53-dependent transcription. **Fig. 8A.** SW480 cells were cotransfected with the PG13-LUC reporter (1 μ g), p53 (2 μ g) and either wt BRCA1 or its mutant derivatives (3.5 μ g) as indicated "+". **Fig. 8B.** SW480 cells were transfected with the -143-LUC p21 reporter (1 μ g) and different combinations (as indicated "+") of BRCA1 or its vector, pCR3 (2 μ g) and p53 or its vector, pCEP4 (2 μ g) along with 1 μ g of

~~-143-LUC p21 reporter. Fig. 8C and Fig. 8D. Inhibitory~~
negative effect of BRCA1 deletion mutant on p53 transcription activity. **Fig. 8C.** SW480 cells were transfected with 0.5 μ g of the PG13-LUC reporter and different combinations (as indicated "+") of pCEP4 or p53

(0.25 μ g) and increasing amounts (lane 3, 2 μ g; lane 4, 4 μ g; and lane 5, 8 μ g) of *BRCA1* Δ 500-1863. The total amount of DNA was maintained constant at 8.75 μ g by adding pCR3 DNA. Fig. 8D, SW480 cells were transfected with 0.5 μ g of the PG13-LUC reporter and different combinations (as indicated "+") of pCEP4 or p53 (0.25 μ g), *BRCA1* Δ 500-1863 (2 μ g) and increasing amounts (lane 4, 4 μ g; lane 5, 6 μ g) of wt *BRCA1*. The total amount of DNA was kept constant at 8.75 μ g by adding pCR3 DNA. Cells were harvested 24 hrs later and luciferase activity was measured as described in Example 2.

DETAILED DESCRIPTION OF THE INVENTION

I. Definitions

Various terms relating to the biological molecules of the present invention are used hereinabove and also throughout the specifications and claims.

With reference to nucleic acid molecules, the term "isolated nucleic acid" is sometimes used. This term, when applied to DNA, refers to a DNA molecule that is separated from sequences with which it is immediately contiguous (in the 5' and 3' directions) in the naturally occurring genome of the organism from which it was derived. For example, the "isolated nucleic acid" may comprise a DNA molecule inserted into a vector, such as a plasmid or virus vector, or integrated into the genomic DNA of a procaryote or eucaryote. An "isolated nucleic acid molecule" may also comprise a cDNA molecule.

With respect to RNA molecules, the term "isolated nucleic acid" primarily refers to an RNA molecule encoded by an isolated DNA molecule as defined above. Alternatively, the term may refer to an RNA molecule that has been sufficiently separated from RNA

molecules with which it would be associated in its natural state (i.e., in cells or tissues), such that it exists in a "substantially pure" form (the term "substantially pure" is defined below).

- 5 With respect to proteins or peptides, the term "isolated protein (or peptide)" or "isolated and purified protein (or peptide)" is sometimes used herein. This term refers primarily to a protein produced by expression of an isolated nucleic acid molecule of the invention.
- 10 Alternatively, this term may refer to a protein which has been sufficiently separated from other proteins with which it would naturally be associated, so as to exist in "substantially pure" form.

- The term "substantially pure" refers to a
- 15 preparation comprising at least 50-60% by weight the compound of interest (e.g., nucleic acid, oligonucleotide, protein, etc.). More preferably, the preparation comprises at least 75% by weight, and most preferably 90-99% by weight, the compound of interest.
- 20 Purity is measured by methods appropriate for the compound of interest (e.g. chromatographic methods, agarose or polyacrylamide gel electrophoresis, HPLC analysis, and the like).

- Nucleic acid sequences and amino acid sequences
- 25 can be compared using computer programs that align the similar sequences of the nucleic or amino acids thus define the differences. For purposes of this invention, the GCG Wisconsin Package version 9.1, available from the Genetics Computer Group in Madison, Wisconsin, and the
-
- 30 default parameters used (gap creation penalty=12, gap extension penalty=4) by that program are the parameters intended to be used herein to compare sequence identity and similarity.

The term "substantially the same" refers to nucleic acid or amino acid sequences having sequence variation that do not materially affect the nature of the protein (i.e. the structure, thermostability characteristics and/or biological activity of the protein). With particular reference to nucleic acid sequences, the term "substantially the same" is intended to refer to the coding region and to conserved sequences governing expression, and refers primarily to degenerate codons encoding the same amino acid, or alternate codons encoding conservative substitute amino acids in the encoded polypeptide. With reference to amino acid sequences, the term "substantially the same" refers generally to conservative substitutions and/or variations in regions of the polypeptide not involved in determination of structure or function.

The terms "percent identical" and "percent similar" are also used herein in comparisons among amino acid and nucleic acid sequences. When referring to amino acid sequences, "percent identical" refers to the percent of the amino acids of the subject amino acid sequence that have been matched to identical amino acids in the compared amino acid sequence by a sequence analysis program. "Percent similar" refers to the percent of the amino acids of the subject amino acid sequence that have been matched to identical or conserved amino acids. Conserved amino acids are those which differ in structure but are similar in physical properties such that the exchange of one for another would not appreciably change the tertiary structure of the resulting protein. Conservative substitutions are defined in Taylor (1986, J. Theor. Biol. 119:205). When referring to nucleic acid molecules, "percent identical" refers to the percent of

the nucleotides of the subject nucleic acid sequence that have been matched to identical nucleotides by a sequence analysis program.

With respect to antibodies, the term
5 "immunologically specific" refers to antibodies that bind to one or more epitopes of a protein of interest, but which do not substantially recognize and bind other molecules in a sample containing a mixed population of antigenic biological molecules.

10 With respect to oligonucleotides or other single-stranded nucleic acid molecules, the term "specifically hybridizing" refers to the association between two single-stranded nucleic acid molecules of sufficiently complementary sequence to permit such
15 hybridization under pre-determined conditions generally used in the art (sometimes termed "substantially complementary"). In particular, the term refers to hybridization of an oligonucleotide with a substantially complementary sequence contained within a single-stranded
20 DNA or RNA molecule, to the substantial exclusion of hybridization of the oligonucleotide with single-stranded nucleic acids of non-complementary sequence.

The term "promoter region" refers to the transcriptional regulatory regions of a gene, which may
25 be found at the 5' or 3' side of the coding region, or within the coding region, or within introns.

The term "selectable marker gene" refers to a gene encoding a product that, when expressed, confers a selectable phenotype such as antibiotic resistance on a
30 transformed cell.

The term "reporter gene" refers to a gene that encodes a product which is easily detectable by standard methods, either directly or indirectly.

The term "operably linked" means that the regulatory sequences necessary for expression of a particular coding sequence are placed in the DNA molecule in the appropriate positions relative to the coding sequence so as to enable expression of the coding sequence. This same definition is sometimes applied to the arrangement of transcription units and other regulatory elements (e.g., enhancers or translation regulatory sequences) in an expression vector.

10

II. Description

A significant proportion of hereditary breast and ovarian cancer predisposition has been attributed to inherited defects in the *BRCA1* tumor suppressor gene.

15 *BRCA1* is a nuclear protein with properties of a transcription factor, and can interact with the recombination and repair protein RAD51. Young women with germline alterations in *BRCA1* develop breast cancer at rates 100-fold higher than the general population and
20 *BRCA1*-null mice die before day 8 of development, but the mechanisms of *BRCA1*-mediated growth regulation and tumor suppression heretofore has remained unknown.

In accordance with the present invention, it has now been shown that *BRCA1* transactivates expression
25 of the CDK-inhibitor *p21^{WAF1/Cip1}* in a p53-independent manner and that *BRCA1* inhibits S-phase cell cycle progression following its transfection into human cancer cells. *BRCA1* does not inhibit S-phase progression in *p21*^{-/-} as compared to *p21*^{+/+} cells and tumor-associated
30 transactivation-deficient mutants of *BRCA1* are defective in both transactivation of *p21* and cell cycle inhibition. Thus, one mechanism by which *BRCA1* contributes to cell cycle arrest and growth suppression is through p53-

independent induction of p21.

Deletion mapping of the human p21 promoter identified a control region of 50 base pairs within the proximal promoter, -143 to -93 of the human p21, that
5 mediates activation of p21 by BRCA1 (Fig. 2C). Deletion mapping further revealed that the two previously described p53 binding sites were not required for BRCA1 transactivation of p21.

The inventors have also discovered that BRCA1
10 increases p53-dependent transcription from the p21 and bax promoters and that BRCA1 and p53 proteins interact both *in vitro* and *in vivo*. The interacting regions map, *in vitro*, to amino acid residues 224-500 of human BRCA1 (SEQ ID NO:1) and the C-terminal domain of p53 (residues
15 300-393 of SEQ ID NO:2). Tumor-derived transactivation deficient BRCA1 mutants are defective in co-activation of p53-dependent transcription and a truncation mutant of BRCA1 that retains the p53-interacting region acts as a dominant inhibitor of p53-dependent transcription. BRCA1
20 and p53 cooperatively induce apoptosis of cancer cells. These results indicate that BRCA1 and p53 coordinately regulate gene expression in their role as tumor suppressors.

The discoveries made in accordance with the
25 present invention can be utilized for a variety of purposes, including but not limited to (1) assay systems to identify pharmacological agents capable of suppressing or enhancing cell proliferation by affecting the p53-
dependent or -independent BRCA1-mediated transcription
30 pathways described above; (2) diagnostic methods for assessing functional mutations in BRCA1 and /or p53 relating to their ability to associate with each other or with p21 regulatory regions or regulatory regions of

other downstream targets involved in cell cycle arrest or apoptosis; and (3) methods for controlling cell proliferation by regulating either cell cycle progression or apoptosis via p53-dependent or -independent BRCA1-mediated transcription pathways, particularly via p21.

The following description sets forth the general procedures involved in practicing the present invention. To the extent that specific materials are mentioned, it is merely for purposes of illustration and is not intended to limit the invention. Unless otherwise specified, general cloning procedures, such as those set forth in Sambrook et al., Molecular Cloning, Cold Spring Harbor Laboratory (1989) (hereinafter "Sambrook et al.") or Ausubel et al. (eds) Current Protocols in Molecular Biology, John Wiley & Sons (1999) (hereinafter "Ausubel et al.") are used.

III. Biological Molecules and Cells

In addition to cells and biological molecules used for general cloning and expression, the following biological molecules and cell types are either required or useful for practice of the methods of the present invention: (1) cloning or expression vectors comprising nucleic acids encoding BRCA1 and p53, or specific portions thereof; (2) reporter genes operably linked to promoters responsive to p53 and/or BRCA1; (3) isolated BRCA1 or p53 protein, or selected portions thereof; (4) antibodies immunologically specific for BRCA1, p53 or selected portions thereof; and (5) various cell lines, including lines deficient in functional p53, BRCA1 (or both), or deficient in downstream targets of p53 or BRCA1, and further engineered with specified reporter genes.

Any nucleic acid molecule that encodes BRCA1 or p53 is contemplated for use in the present invention. Likewise, any BRCA1 or p53 protein is considered suitable for use in the present invention. Human BRCA1 and p53 are exemplified herein (e.g., SEQ ID NO:1 and SEQ ID NO:2). Another preferred source of BRCA1 and p53 proteins and nucleic acids is the mouse. Any other BRCA1 and p53 nucleic acids also may be used. These genes and proteins have been extensively characterized, and sequence information is widely available, e.g., through GenBank and the published and patent literature. For instance, U.S. Patent No. 5,654,155 discloses a consensus DNA sequence for the *BRCA1* gene. U.S. Patent 5,750,400 discloses three *BRCA1* coding sequences; the following Genbank Accession Numbers contain *BRCA1* protein sequences: 1147603, 1418288, 3994048, 3994047, 3994046, 3209867, 2489823, 1698399, 728984, 627392, 1620568.

Included in the present invention is a nucleic acid molecule comprising the region of the proximal promoter of the *p21* gene responsible for p53-independent *BRCA1* transactivation of *p21*. Although the *BRCA1*-responsive element from *Homo sapiens* is described and exemplified herein (-143 to -93 from the transcription start site noted in SEQ ID NO:3), this invention is intended to encompass the *BRCA1* response element of the *p21* promoter from other mammals that are sufficiently similar to be used instead of the *Homo sapiens* nucleic acid sequence for *BRCA1*-mediated transactivation. These include, but are not limited to, allelic variants and natural mutants in the relevant region of SEQ ID NO:3 (bases 4441-4491), as well as homologs found in different species of mammals. Because such variants are expected to possess certain differences in nucleotide sequence,

this invention provides an isolated nucleic acid molecule having at least 60% (preferably 70% and more preferably over 80%) sequence homology to the nucleotide sequence set forth as residue -143 extending to the transcription start site noted in SEQ ID NO:3 over the length of those residues (residues -143 to -93 comprise the specific element conferring transactivation by BRCA1; however, the intervening 5' promoter region comprises regulatory elements needed for transcription (for instance, the TATA box). Because of the natural sequence variation likely to exist among BRCA1 transactivated genes, one skilled in the art would expect to find up to about 30-40% nucleotide sequence variation, while still maintaining the unique properties of the BRCA1 response element of the human p21 promoter.

Peptides comprising the regions of BRCA1 and p53 constituting the protein-protein binding site, and nucleic acids encoding those peptides, are included in the present invention. As binding sites do not require absolute conservation of sequence and a some substitution of amino acids of similar chemical characteristics is allowed, the invention is contemplated to include peptides comprising sequences similar residues 224 to 500 of SEQ ID NO:1 or residues 300-393 of SEQ ID NO:2. In a preferred embodiment, peptides are 80% similar to residues 224-500 of SEQ ID NO:1 or residues 300 to 393 of SEQ ID NO:2 over the length of aa residues 224 to 500 of SEQ ID NO:1 or residues 300-393 of SEQ ID NO:2; (more preferably peptides are 90% similar; and most preferably peptides are 95% similar).

The use of the p53-binding sites to confer or remove p53 activation and/or p53/BCRA1 coactivation upon promoters is considered to be part of the invention. The

removal of the p53 binding sites from p53-regulated promoters both *in vitro* and *in vivo* will allow these promoters to be used in constructs without p53 regulation as well as to create cell lines with altered p53 regulation. Promoters without p53 regulation may be changed to p53 regulation through the inclusion of p53-binding sites. It is further contemplated that variations of the p53-binding site sequence may be used for this invention.

10 The use of BCRA1 sequence without the transactivation domain but with the p53 binding domain on BCRA1 to lessen p53 regulation cell proliferation is considered part of the invention. In a preferred embodiment, the region of BCRA1 with the p53-binding domain and without the transactivation domain is used. In accordance with this aspect of the invention it is now known that the BCRA1/p53 interaction will be hindered by the addition of partially functional binding domains, so it is highly likely that the BCRA1 binding domain on p53 will likewise repress the BCRA1/p53 coactivation of genes. In another preferred embodiment, the region of p53 with the BCRA1 binding domain (residues 300 to 393 of SEQ ID NO:2) without the DNA-binding domain is used. In a more preferred embodiment, the p53 binding domain on BCRA1 (residues 224 to 500 of SEQ ID NO:1) or the BCRA1 binding domain on p53 (residues 300-393 of SEQ ID NO:2) is used. In a most preferred embodiment, the truncated BCRA1, BCRA1 Δ 500-1863 (residues 1 to 499 of SEQ ID NO:1), is used. It is contemplated that proteins significantly similar to these proteins can also be used in this invention. In a preferred embodiment, proteins 60% similar to residues 1-499 of SEQ ID NO:1 are used, in a more preferred embodiment proteins 70% similar to

residues 1-499 of SEQ ID NO:1 are used, and in a most preferred embodiment, proteins 80% similar to residues 1-499 of SEQ ID NO:1 are used.

Nucleic acid molecules used in the invention
5 may be prepared by two general methods: (1) they may be synthesized from appropriate nucleotide triphosphates, or (2) they may be isolated from biological sources. Both methods utilize protocols well known in the art. In addition, some of the nucleic acid molecules may be
10 commercially available.

Proteins used in the present invention may be prepared in a variety of ways, according to known methods. The protein may be purified from appropriate sources, e.g., human or animal cultured cells or tissues,
15 by immunoaffinity purification. However, this is not a preferred method due to the low amount of protein likely to be present in a given cell type at any time.

The availability of nucleic acids molecules encoding BRCA1, p53 and portions thereof enables
20 production of the protein using *in vitro* expression methods known in the art. For example, a cDNA or gene may be cloned into an appropriate *in vitro* transcription vector, such a pSP64 or pSP65 for *in vitro* transcription, followed by cell-free translation in a suitable cell-free
25 translation system, such as wheat germ or rabbit reticulocytes. *In vitro* transcription and translation systems are commercially available, e.g., from Promega Biotech, Madison, Wisconsin or BRL, Rockville, Maryland.

Alternatively, according to a preferred
30 embodiment, larger quantities of protein may be produced by expression in a suitable procaryotic or eucaryotic system. For example, part or all of a DNA molecule may be inserted into a plasmid vector adapted for expression in a bacterial cell, such as *E. coli*, or into a
35 baculovirus vector for expression in an insect cell.

Such vectors provide the regulatory elements necessary for expression of the DNA in the given host cell, positioned in such a manner as to permit expression of the DNA in the host cell. Such regulatory elements
5 required for expression include promoter sequences, transcription initiation sequences and, optionally, enhancer sequences.

Proteins produced by gene expression in a recombinant procaryotic or eucaryotic system may be
10 purified according to methods known in the art. In a preferred embodiment, a commercially available expression/secretion system can be used, whereby the recombinant protein is expressed and thereafter secreted from the host cell, to be easily purified from the
15 surrounding medium. If expression/secretion vectors are not used, an alternative approach involves purifying the recombinant protein by affinity separation, such as by immunological interaction with antibodies that bind specifically to the recombinant protein. Such methods
20 are commonly used by skilled practitioners.

Some aspects of the present invention also utilize antibodies which bind immunospecifically to BRCA1 or P53 epitopes or proteins. Such antibodies may be prepared according to standard methods, if they are not
25 commercially available. In a preferred embodiment, monoclonal antibodies are prepared, which react immunospecifically with various epitopes of BRCA1 or p53. Monoclonal antibodies may be prepared according to general methods of Köhler and Milstein, following
30 standard protocols. Polyclonal or monoclonal antibodies that immunospecifically interact with BRCA1 or p53 can be utilized for purifying the proteins, according to standard methods. They also may be used in the diagnostic and therapeutic methods described below.

35 Vectors that express the BRCA1 and p53

proteins, protein fragments, mutants, etc. are used in a variety of assays of the invention, as described below. For instance, the examples teach the transfection of cells with pCEP4-p53 and pCR3-BRCA1 expression vectors.

5 Any vector that expresses the desired coding sequence in adequate amounts in the cell line of choice is acceptable. If non-*Homo sapiens* cell lines are used, different expression vectors may be required to achieve adequate expression of the protein, as is well known in

10 the art. Many different kinds of expression vectors are commercially available (e.g., from Invitrogen).

Reporter genes operably linked to BRCA1- or p53-responsive promoters are also used in certain of the assays described below. Examples of several such

15 reporter plasmids specific to p53/BRCA1 activation are taught in Examples 1 and 2. PG13-LUC, pbax-LUC and pWWP-LUC are all reporter plasmids that express luciferase. The use of the luciferase as a reporter molecule is well known in the art. The aforementioned reporter plasmids

20 use three different promoters that are all coactivated by BRCA1/p53. The p21 promoter of the pWWP-LUC reporter plasmid alone exhibits p53-dependent and p53-independent coactivation. The use of pWWP-LUC plasmid is therefore particularly suited to address how a test molecule may

25 differentially affect these two aspects of BRCA1 activation. Other reporter genes may be expressed, and many are commercially available. Reporter genes of interest include, but are not limited to, those encoding green fluorescent proteins (GFPs), β -glucuronidase (GUS)

30 and β -galactosidase, among others.

A variety of cell types can be used to practice different aspects of the present invention. In the assays described below for screening potential

pharmacological agents, cell lines deficient in one or more of BRCA1, p53 or transactivational targets of BRCA1 and/or p53, are preferred. For instance, Example 2 teaches that SW480 human colon adenocarcinoma cells and mouse embryo fibroblast cells may be used. In general, tumor cells will provide the best source of regulation-deficient lines for use in the present invention. However, as discussed below, normal cells may also be used. For instance, to expand a normal cell line, the BRCA1-mediated transactivation of tumor suppressor genes would be down-regulated in order to stimulate proliferation of the cells.

Cells lines additionally do not need to be from humans or mice, but may be from other mammalian sources, such as rats, pigs, dogs, cows and primates, to name a few. Other useful cell lines and systems for these assays include, but are not limited to, cell lines derived from *Drosophila melanogaster* or *Caenorhabditis elegans*.

IV. Methods and Kits

Provided in accordance with the present invention are assay systems that can be used to discover and develop pharmacological agents capable of affecting the regulation of BRCA1-activated gene transcription, and their use in the treatment and control of cell proliferative disorders. The assay systems can be used in their entirety in cells, or partially *in vitro*.

One class of assays of the invention are assays for molecules that participate in the BRCA1/p53 cell proliferation regulation pathway. These assays use the BRCA1/p53 interaction as a basis for determining the degree of specific physical interaction that a test

molecule has with p53 and/or BRCA1, and the effect of the test molecule will have on the cell cycle progression. These assays may be performed *in vitro* and *in vivo*.

One assay of this type is an *in vitro* assay
5 that determines the effect of test molecules on the protein-protein interaction between p53 and BRCA1. While this assay has many embodiments, the basic assay consists of three steps:

1. Incubating BRCA1 and p53 in the presence or
10 absence of the test molecule;

2. Separating complexes comprising p53 and BRCA1 from the unbound molecules; and

3. Determining the effect of the test molecule on the formation of complexes comprising p53 and BRCA1.

15 Each of these step has many variations, the choice of which will customize the assay for a particular use.

In step 1, BRCA1 and p53 may be provided as isolated proteins (e.g., produced separately by *in vitro*
20 transcription/translation or by *in vivo* expression). Alternatively, vectors that express genes encoding the proteins may be utilized. The test compound can be any of a variety of substances, including but not limited to small molecules, peptides, proteins, nucleic acids, or
25 any combination thereof.

In step 2, numerous variations exist, the choice of which depends on the laboratory facilities and skills available. Example 2 teaches the use of
glutathione-sepharose beads to precipitate glutathione S-
30 transferase (GST) fusion proteins. Example 2 also teaches the use of antibodies specific to p53 or BRCA1 to precipitate complexes. Many variations exist on these two methods. Kits are commercially available with

different kinds of beads and methods to bind to proteins of interest. These beads may be separated from the total assay buffer by centrifugation, filtering, or in the case of magnetic beads, magnetism. Numerous methods exist
5 that will be well known and usable by those skilled in the art.

Step three determines the molecules that are in the isolated complexes. If a non-protein test molecule has been employed in step one, the outcome in step three
10 may be that BRCA1 does not associated with p53 as well as the controls, or BRCA1 may associate with p53 better than the controls. In Example 2, SDS-PAGE is taught as a method for identifying the components of the complexes by molecular weight. If the protein molecules of the assay
15 are labeled with [³⁵S]-methionine as is taught in Example 1, then they can be identified by fluorography. [³⁵S]-methionine labeling additionally allows the amount of the protein to be accurately determined and allows the detection of levels of protein too low for Coomassie
20 staining and silver staining after SDS-PAGE. Additionally, proteins in the precipitated complex can be identified by western blotting after PAGE according to methods well known to those skilled in the art. Anti-BRCA1 and anti-p53 monoclonal antibodies are commercially
25 available (Oncogene Science).

Another assay of the invention is a cell-based assay to determine the effect of a particular test molecule on p53-dependent and p53-independent BRCA1
regulation of cell proliferation regulation pathways. In
30 accordance with the present invention it has been learned that BRCA1 regulates the cell cycle through p53-dependent coactivation and p53-independent activation of genes. While this assay has many embodiments, the basic assay

consists of two steps:

1. Cotransfection of a cell line that lacks functional p53 and BRCA1 with a series of vector combinations that express p53, BRCA1, and empty vector controls, for instance, such as those shown in the following array, along with a reporter gene operably linked to a promoter activated by p53 and/or BRCA1:

10	BRCA1 expression vector	-	+	-	+	-	+	-	+
	empty BRCA1 expression vector	+	-	+	-	+	-	+	-
15	p53 expression vector	-	-	+	+	-	-	+	+
20	empty p53 expression vector	+	+	-	-	+	+	-	-
	Test molecule	-	-	-	-	+	+	+	+
25									

2. Determine the level of activity of the BRCA1/p53 cell proliferation pathway.

Each of these steps has numerous variations, the choice of which will customize the assay to specific conditions. This assay can be used to particular advantage in conjunction with the *in vitro* assay. Test molecules that are found to inhibit the p53/BRCA1 interaction in the *in vitro* assay can be tested for their *in vivo* action in this cell-based assay.

In the first step, a cell line that is deficient in p53 and/or BRCA1 is used, and may be further engineered with reporter genes operably linked to p53-specific or BRCA1-specific promoters, as described for instance in Example 2. The transfection methods taught in Example 2 are lipofectin, lipofectamine, and calcium

phosphate-precipitation. The choice of transfection method depends on the type of cell used in the transfection, and such methods are well known to persons skilled in the art. Additionally, cell lines stably
5 transformed with the appropriate expression cassettes may be used instead of transfection.

Step two of the cell based assay requires that the activity of the BRCA1/p53 cell proliferation pathway be determined. Two general methods are taught in the
10 examples. Reporter genes may be co-transfected into cells. Alternatively, an apoptosis assay or cell cycle arrest assay may be used as a means of determining the activity of the BRCA1 or BRCA1/p53 cell proliferation pathways. The TUNEL assay is taught in Example 2 as a
15 method for determining the degree of cell apoptosis. This method, and other methods that measure the gross outcome of BRCA1/p53 cell proliferation pathway regulation, are very useful for determining the therapeutic and industrial usefulness of test compounds.
20 Other methods that measure this broad scope of effect include, but are not limited to, the BrdU incorporation assay to measure DNA synthesis taught in Example 1 and assays for activity of apoptosis-related enzymes, e.g., caspase. Various assays to measure cell proliferation,
25 cell cycle arrest and apoptosis are known to persons skilled in the art.

The present invention also includes diagnostic assays. For instance, standard genetic screening for mutations in the p53/BRCA1 respective binding domains, or
30 the domain responsible for p53-independent binding of BRCA1 to the p21 promoter, is now possible in accordance with the present invention. Such screening assays may utilize the specific nucleic acid molecules as

hybridization probes. Alternatively, antibodies immunologically specific for the respective binding domains may be developed and utilized.

Nucleic acid molecules encoding BRCA1 and p53
5 (and specific portions thereof, as well as part or all of the proteins themselves, may be used to advantage to control cellular proliferation, either through p53-independent regulation of p21 or through co-regulation of apoptosis signal transduction by BRCA1 and p53. In a
10 preferred embodiment, the p53-independent BRCA1 activation pathway or the BRCA1/p53 co-activation pathway is up-regulated, thereby inducing cell cycle arrest or apoptosis. Such up-regulation is suitable for treatment of neoplastic cell growth either in cultured cells or in
15 vivo, in a patient requiring such therapy.

In another embodiment, the aforementioned pathways are down-regulated, resulting in an increase in cell proliferation due to the removal of cell cycle checkpoints and apoptosis signal transduction. This type
20 of down regulation is useful for expanding cell lines in culture, or as a research tool to study the effect of such manipulation on long term cell growth and development.

Up-regulation or down-regulation of p53-
25 independent or -dependent BRCA1-mediated transcription may be accomplished in a variety of ways known to persons skilled in the art. Some of these ways are described above, and utilize various DNA or protein fragments, or
truncation mutants of BRCA1, to block the pathway(s) at
30 various points, thereby down-regulating the pathway(s) and increasing cell proliferation. Other methods include, but are not limited to, (1) over-producing BRCA1 or p53 in cells that do not produce the proteins, or that

produce defective proteins; and (2) enhancing the interaction between BRCA1 and p53 or the p21 promoter. Such methods are considered within the scope of the present invention.

5 According to another aspect of the invention, kits are provided to facilitate performing the above-described assays. In one embodiment, the kits comprise one or more of the DNA constructs encoding BRCA1, p53, fragments and mutants thereof, and reporter genes
10 operably linked to transactivational targets of BRCA1 and/or p53, as described in greater detail herein, along with instructions on how to use the constructs to perform the assays of the invention. In another embodiment, the kits comprise aliquots of transgenic cells and
15 instructions for their use. In another embodiment, the kits may comprise antibodies and other reagents for performing immunological assays. The kits may also comprise, optionally, various additional reagents for the assays, such as growth media, enzyme substrates for the
20 reporter gene product, and standard solutions for calibrating expression of the reporter gene.

The following examples are provided to describe the invention in greater detail. They are intended to illustrate, not to limit, the invention.

25

EXAMPLE 1

30 BRCA1 Arrests the Cell Cycle by
Transactivating the Expression of p21^{WAF1/CIP1}

Because several known tumor suppressor genes interact with or negatively regulate the cell cycle machinery, the effect of BRCA1 on cell cycle progression
35 was investigated. This example describes the results of that investigation.

A. Materials and Methods

Plasmid constructs. The pCEP4-p53 and pWWP-Luc plasmids are described in the literature (El-Deiry et al., 1993, *supra*). The construction of the p21-promoter deletions fused to the luciferase reporter gene has been described (Zeng et al., 1997, *Nature Genetics* 15:78-82). The human p21-promoter luciferase-reporters with 5'-truncations at -153, -143 and -133 were constructed and sequenced as described (Zeng et al., 1997, *Nature Genetics* 15:78-82). The murine p21 promoter-CAT reporter, pCAT1, has been described (El-Deiry et al., 1995, *Cancer Res.* 55:2910-2919). The GFP expression vector pGreen Lantern-1 was obtained from Gibco. The pCR3-BRCA1 expression plasmid has been previously described (Thankur et al., 1997, *Mol. Cell. Biol.* 17:4444-452).

pCR3 vectors encoding synthetic and tumor-associated BRCA1 mutants (Fig. 3A) were constructed as follows. The RAD51-interaction-deficient mutant plasmid (Δ 515-1091) was constructed by digestion of pCR3-BRCA1 using *Bsu36I* to delete nucleotides 1661-3392 (Miki et al., 1994, *Science* 266:66-71), followed by in-frame intramolecular ligation. The transactivation domain deletion mutant plasmid (Δ 1314-1863) was prepared by digestion of pCR3-BRCA1 with *Bam*H1 (nucleotide 4058) and *Not*I (nucleotide 5833), Klenow fill-in and subsequent intramolecular blunt-end ligation at 37°C. The double mutant (Δ 500-1863) was cloned by polymerase chain reaction (PCR) amplification of nucleotides 1-1616 using pCR3-BRCA1 as a template and the following primers:

5'-GCAAGCTTGCCACCATGGATTTATCTGCTCTTCGC-3' (SEQ ID NO:4) and

5'-TTGTGAGGGGACGCTCTTGTA-3' (SEQ ID NO:5).

The 1.6-kilobase PCR product was ligated into the pCR3 vector, and a clone expressing the N-terminal region of BRCA1 in the sense orientation downstream of the CMV-promoter was isolated. The Δ NLS/C + NLS vector was
5 prepared as follows. A 488-bp DNA fragment containing a fusion between the extreme C terminus of BRCA1 and the NLS region (amino acids 499-510) was generated by PCR amplification of pCR3-BRCA1 using the following primers:

5'-AGGAGATGTGGTCAATGGAAG-3' (SEQ ID NO:6) and
10 5'-TATCATGATGTAGGTCTCCTTTTACGCTTTAATTATT
GTAGTGGCTGTG-3' (SEQ ID NO:7).

This PCR product was subcloned into pCR3 and released as an *Apal* fragment which was cloned in-frame into an *Apal*-digested Δ NLS plasmid. pCR3 vectors encoding the
15 transactivation-domain tumor-derived BRCA1 mutants (Chapman and Verma, 1996, *supra*; Monteiro et al., 1996, *supra*) (P1749R, Y1853insA, Q1756insC) were constructed by amplification of a 1.8-kb C-terminal region containing the specific mutations and subcloning into pCR3-BRCA1
20 digested with *Bam*H1 and *Not*I. All mutant BRCA1 expression plasmids used were sequenced and shown to express protein. Deletion of the nuclear localization signal (Thakur et al., 1997, *supra*) (Δ NLS mutant) resulted in cytoplasmic staining, which reverted back to
25 the nucleus upon addition of a C-terminal NLS (Δ NLS/C + NLS; unpublished data).

Cells, transfections and luciferase assays.

The p21^{+/+} parental and p21^{-/-} HCT116 human colon cancer cells (Waldman et al., 1995, *Cancer Res.* 55:5187-5190)
30 were provided by T. Waldman and B. Vogelstein. SW480, HeLa, COS-7 and CV1 cells (ATCC) were transfected as described (El-Deiry et al., 1993, *Cell* 75:817-825). Luciferase and CAT assays were performed as described

(El-Deiry et al., 1993, *supra*; El-Deiry et al., 1995, Cancer Res. 55:2910-2919). HeLa cells expressing BRCA1 or mutants were obtained following transfection of HeLa cells with BRCA1, Δ NLS or Δ 515-1091 mutants and
5 continuous growth in $0.4 \text{ mg} \cdot \text{ml}^{-1}$ G418.

Analysis of BrdU incorporation in transfected GFP(+) cells. SW480 or HCT116 cells were co-transfected with pGreen Lantern-1 and mammalian expression vectors (as indicated in Figs 1,4 and Table 1) at a ratio of 1:3.
10 At 12 h after transfection, BrdU (Sigma) was added at a final concentration of $20 \mu\text{M}$ and the cells incubated for 20 h at 37°C . Cells were examined by fluorescence microscopy to identify GFP(+) cells. Cells were treated with a mouse anti-BrdU monoclonal antibody (BM 9318,
15 Boehringer Mannheim) and Rhodamine-conjugated goat anti-mouse IgG (Pierce) as described (Zeng et al., 1997, *supra*). The number of BrdU(+) cells was determined for all of the GFP(+) cells in three high-power fields, as described in Fig. 1.

Immunocytochemistry and immunofluorescence.
SW480 cells, transfected with expression plasmids, were stained 24 h later with an anti-human-WAF1 monoclonal antibody (Ab1; Calbiochem) as described (El-Deiry et al., 1995, *supra*). Immunofluorescence analysis of BRCA1
25 expression was performed as described (Thakur et al., 1997, *supra*).

Northern blot analysis. Total RNA was isolated and northern blot analysis was performed as described (El-Deiry et al., 1993, *supra*). and p21-mRNA-expression
30 was detected using a 2.1-kb human p21 cDNA probe (El-Deiry et al., 1993, *supra*). Equivalent loading of various RNA samples was demonstrated using a probe for rpl32, which encodes a ribosomal protein (Meyuhas and

Perry, 1980, Gene 10:113-129).

Immunoprecipitation and western blot analysis.

Cells were lysed in a buffer containing 50 mM Tris-HCl, pH 7.5, 120 mM NaCl, 50 mM NaF, 0.5% NP-40, 1 mM EDTA, pH 8.0, 1 mM phenylmethylsulphonyl fluoride (Gibco), 1% antipain (Sigma), 1% leupeptin (Sigma), 1% pepstatin A (Sigma), 1% chymostatin (Sigma) and 1% AEBSF (Calbiochem). Immunoprecipitations were carried out in the lysis buffer using 2 µg of anti-BRCA1 monoclonal antibody (Ab1; Calbiochem) for 2 h at 4°C, followed by the addition of 50% protein A-agarose beads (Sigma) and incubation for 1 h. After 3 washes with lysis buffer the immunoprecipitated proteins were analyzed by western blotting as described in Example 1, using a 1:250 dilution of the anti-BRCA1 monoclonal antibody, which was raised using the N-terminal portion of recombinant human BRCA1 (amino acids 1-304) as the immunogen (Calbiochem).

B. Results

By using green fluorescent protein (GFP) to mark specific transfected cells (Zeng et al., 1997, Biotechniques 23:88-94), we examined the effects of BRCA1 transfection on new DNA synthesis in SW480 and HCT116 human colon cancer cells (Figs 1, 4 and Table 1). We found fewer BrdU(+)/GFP(+) SW480 cells following transfection of either BRCA1 or p53 than with their control vectors (Fig. 1). A quantitative summary of the percentage of BrdU(+)/GFP(+) cells from three independent experiments in SW480 cells is shown in Fig. 1 and Table 1. BRCA1 inhibited new DNA synthesis in SW480 cells by approximately 50% compared with the pCR3 vector. The extent of inhibition of BrdU incorporation following BRCA1 transfection was similar to p53 transfection (Fig. 1). BRCA1 also inhibited S-phase progression in HCT116

cells (Fig. 4). These results suggest that BRCA1 can inhibit S-phase progression and thus negatively regulate the cell cycle in human cancer cells.

5 **Table 1.** Cell cycle effects of tumor-derived BRCA1 mutants.

		GFP(+) cells	BrdU(+) cells	BrdU(+)/GFP(+) (%)
10	pCR3	43	27	62.8
	BRCA1	55	19	34.6
	P1749R	59	31	52.5
15	Y1853insA	56	32	57.1

20 These BRCA1 mutants are defective at inhibiting DNA synthesis. GFP was used as a marker for transfection of SW480 cells. GFP(+) cells were examined for BrdU incorporation (BrdU(+)) by anti-BrdU staining. BrdU(+)/GFP(+) cells with active DNA synthesis.

We investigated cyclin-dependent kinase (CDK) inhibition of cell-cycle progression as a potential mechanism by which BRCA1 may control cell proliferation.

25 Induction of p21^{WAF1/CIP1} expression has been linked to growth inhibition by p53 (El-Deiry et al., 1993, Cell 75:817-825)), and p21 expression also has been found to signal growth arrest, independent of p53, in cells undergoing differentiation (Zhang et al., 1995, Cancer

30 Res. 55:668-674). The protein p21 is a universal cell-cycle inhibitor that specifically binds cyclin-CDK complexes and proliferating cell nuclear antigen, thereby serving as a potent growth inhibitor and effector of cell-cycle checkpoints (Sherr and Roberts, 1995, Genes

35 Dev. 9:1149-1163). As BRCA1 contains a carboxy-terminal transactivation domain (Chapman and Verma, 1996, Nature 382:678-679), we hypothesized that BRCA1 may transcriptionally induce p21 expression and thus

negatively regulate cell-cycle progression. We examined the effect of BRCA1 on p21-promoter reporter gene expression following transfection into SW480, HCT116, COS-7, HeLa and CV1 cells (Figs 2, 3, and data not shown). BRCA1 activated the human p21 promoter luciferase-reporter by 5- to greater than 20- fold in SW480 (Fig. 2A), HCT116 (Fig. 2A), HeLa (data not shown) and COS-7 (Fig. 3B) cells, as compared to transfection of the pCR3 vector. BRCA1 also transactivated the mouse p21-promoter by more than 10-fold in CV1 cells (Fig. 2B).

Deletion mapping within the human p21 promoter identified a control region of 50 base pairs (between -143 and -93) within the proximal promoter that seems to mediate activation of p21 by BRCA1 (Fig. 2C). The two p53-binding sites are not required for BRCA1 transactivation of p21. Whether p21 activation by BRCA1 is a direct consequence of BRCA1 binding to the p21 promoter or is an indirect effect is not known. We also investigated whether BRCA1 could activate endogenous p21 mRNA and protein expression. Figure 3D shows that p21 mRNA levels were elevated in HeLa cells after BRCA1 transfection. By using immunochemical methods, we also found increased levels of endogenous p21 protein in SW 480 cells transfected with BRCA1, compared with cells transfected with vector alone. Thus, transcriptional activation of p21 by BRCA1 appears to be functionally relevant.

To further elucidate the biological importance of p21 regulation by BRCA1, we examined the effect of various synthetic and tumor-associated mutant BRCA1 proteins on p21 expression and cell-cycle progression (Figs. 3,4 and Table 1). Mutants of BRCA1 lacking a functional nuclear localization signal, the C-terminal

transactivation domain, the RAD51-interacting domain or all three domains were deficient in activating p21 expression (Fig. 3A,B). Similarly, three different tumor-associated transactivation-deficient (Chapman and
5 Verma, 1996, *supra*; Monteiro et al., 1996, PNAS 93:13595-13599) BRCA1 mutants were defective in activating the human p21-promoter luciferase-reporter gene (Fig. 3C). The two tumor-associated transactivation-deficient BRCA1 mutants tested for cell-cycle inhibition were also found
10 to be deficient in cell-cycle inhibition in SW480 cells (Table 1). These results indicate that transactivation by BRCA1 may be required for its cell-cycle inhibitory effect, and that tumor-derived BRCA1 mutants may be defective in cell-cycle inhibition.

15 To determine whether p21 is required for the cell-cycle inhibitory effect of BRCA1, we examined the extent of new DNA synthesis following BRCA1 transfection into p21^{+/+} and p21^{-/-} HCT116 cells. BRCA1 inhibited new DNA synthesis in p21^{+/+} HCT116 cells, but there was no
20 evidence of DNA synthesis inhibition resulting from BRCA1 in the p21^{-/-} cells (Fig. 4). These observations indicate that p21 induction may be required for cell-cycle inhibition by BRCA1 in HCT116 cells. Expression of p21 has previously been shown to be required for cell-cycle
25 arrest following γ -irradiation of these cells (Waldman et al., 1995, Cancer Res. 55:5187-5190).

These results demonstrate that BRCA1 can negatively regulate the mammalian cell cycle, and suggest
~~that this effect is at least partly mediated by the~~
30 ability of BRCA1 to induce p21. Although previous studies have reported that expression of BRCA1 is cell-cycle dependent (Vaughn et al., 1996, Cell Growth Differ. 7:711-715; Gudas et al., 1996, 7:717-723; Chen et al.,

1996, Cancer Res. 56:3168-3172), our results demonstrate that BRCA1 can inhibit cell-cycle progression. The absence of this inhibition in p21^{-/-} cells indicates that p21 expression maybe essential for BRCA1 to inhibit new
5 DNA synthesis. Decreased cell-cycle inhibition by transactivation-deficient tumor-derived BRCA1 mutants is consistent with the idea that regulation of p21 by BRCA1 may contribute to growth control. In support of this, recent observations in the yeast *Saccharomyces cerevesiae*
10 demonstrate that the C-terminal 303 amino acids of BRCA1 are sufficient to inhibit yeast colony formation, and that tumor-associated mutations in the context of the 303 amino-acid region fail to inhibit colony growth (Humphrey et al., 1997, PNAS 94:5820-5825).

15 The results reported in this example are at odds with recent observations that cells from BRCA1-null mouse embryos have increased levels of p21 mRNA, which suggest that BRCA1 may suppress p21 expression during development to allow cell growth (Haken et al., 1996, Cell 85:1009-1023). However, p21 protein level and its
20 effect on the cell cycle have not been determined in BRCA1-null embryos, and the mechanism of increased p21 expression remains unclear (Haken et al., 1996, supra). BRCA1 may serve different functions during development
25 and adulthood. It also is possible that the absence of BRCA1 in these cells perturbs a feedback loop controlling expression of p21. Although our data do not provide a clear explanation for this difference, our results
~~demonstrate that BRCA1 can transcriptionally induce p21~~
30 expression and negatively regulate the cell cycle. The identification of BRCA1 as an RNA polymerase II holoenzyme-associated protein provides additional evidence for the role of BRCA1 in transcriptional

activation (Scully et al., 1997, PNAS 94:5605-5610). The importance of this role in tumor suppression is further supported by the fact that about 90% of the mutations in *BRCA1* result in C-terminal truncations that involve the transactivation domain. The loss of cell-cycle inhibition in *p21*^{-/-} cancer cells and the deficiency in *p21* activation and cell-cycle inhibition by tumor-derived *BRCA1* mutants supports the notion that *p21* expression may lead to a quiescent or growth-inhibited state, which may contribute to *BRCA1*-dependent tumor suppression.

EXAMPLE 2

BRCA1 Interacts with *p53* to Regulate Transcription

In this example, the participation *BRCA1* and *p53* may participate in a common pathway of growth regulation is demonstrated. *BRCA1* and *p53* are shown to form stable complexes, both *in vitro* and *in vivo*. *BRCA1* is shown to be a potent coactivator of *p53*-dependent transcription of the *p21* and *bax* genes.

A. Materials and Methods

Cell Lines and Culture Conditions. The SW480 human colon adenocarcinoma cell line was maintained in culture as described in Example 1. HCT116 human colon cancer cells (wt *p53*) were provided by Bert Vogelstein (Johns Hopkins University), and cultured as previously described (McDonald et al., 1996, Cancer Res. 56:2250-2255). Mouse embryo fibroblasts (*p53*^{-/-}) were provided by Tyler Jacks (Massachusetts Institute of Technology). SAOS-2, Hela, COS7, HBL100, U2OS and 293 cells were obtained from ATCC.

Reporters and Expression Plasmids. The pWWP-LUC, PG13-LUC and pCEP4-*p53* plasmids were provided

by B. Vogelstein (Johns Hopkins University). pbax-LUC was obtained from Karen Vousden (NCI, Frederick, MD). The -143 p21 Luc reporter was described in Example 1. pGEX-5X1 was purchased from Pharmacia, and the pCR3 and pCR3.1 were purchased from Invitrogen. The Luciferase T7 control DNA was obtained from Promega. Wild-type and mutant BRCA1 (Δ exon11, Δ 500-1863, Δ 1312-1863, P1749R, Y1853insA and Q1756insC) were described previously (Thakur et al., 1997, *supra*; Somaundaram et al., 1997, *supra*). The p53 cDNA was amplified using the primers 5'-AAGCTTGCCACCATGGAGGAGCCGAGTCA-3' (SEQ ID NO:8) and 5'-ATGCGGCCGCTCAGTCTGAGTCAG-3' (SEQ ID NO:9) and the human p53 cDNA as template. The resulting PCR product was sub-cloned into pCR3.1. Using the human BRCA1 cDNA as template, venous BRCA1 fragments were amplified and cloned into pCR3.1 for *in vitro* translation. The primers used were:

For BRCA1 exon 11 (224-1365):

5'-TAAGCTTGCCACCATGGCTGCTTGTGAATTT-3' (SEQ ID NO:10) and 5'-ACTCGAGTCATAAGTTTGAATCCAT-3' (SEQ ID NO:11);

BRCA1 (1-1056):

5'-GCAAGCTTGCCACCATGGATTTATCTGCTCTTCGC-3' (SEQ ID NO:12) and 5'-GGAGCCCACTTCATTAGTAC-3' (SEQ ID NO:13);

BRCA1(1-98):

5'-GCAAGCTTGCCACCATGGATTTATCTGCTCTTCGC-3' (SEQ ID NO:14) and 5'-ACCTGTGTCAAGCTGAAAAGCAC-3' (SEQ ID NO:15);

BRCA1 (91-500):

5'-AAGCTTGCCACCATGTGTGCTTTTCAGCTT-3' (SEQ ID NO:16) and 5'-TCTCGAGTCAATTTGTGAGGGGACG-3' (SEQ ID NO:17);

BRCA1(224-500):

5'-TAAGCTTGCCACCATGGCTGCTTGTGAATTT-3' (SEQ ID NO:18) and 5'-TCTCGAGTCAATTTGTGAGGGGACG-3' (SEQ ID NO:19).

For production of GST-p53 fusion proteins,

full-length and different fragments of *p53* were amplified using the following primers:

wt *p53*:

5'-GTCGAATTCGAGGAGCCGAGTCAGAT-3' (SEQ ID NO:20) and

5'-ATGCGGCCGCTCAGTCTGAGTCAG-3' (SEQ ID NO:21);

p53 (1-43):

5'-GTCGAATTCGAGGAGCCGAGTCAGAT-3' (SEQ ID NO:22) and

5'-ATGCGGCCGCCAAATCATCCATTGC-3' (SEQ ID NO:23);

p53 (1-200):

5'-GTCGAATTCGAGGAGCCGAGTCAGAT-3' (SEQ ID NO:24) and

5'-TACTCGAGATTTCCTTCCACTCGGAT-3' (SEQ ID NO:25);

p53 (100-300):

5'-ATGAATTCCAGAAAACCTACCAG-3' (SEQ ID NO:26) and

5'-TACTCGAGGGCAGCTCGTGGTG-3' (SEQ ID NO:27);

p53 (201-393):

5'-ATGAATTCTTGCGTGTGGAGTATTTG-3' (SEQ ID NO:28) and

5'-ATGCGGCCGCTCAGTCTGAGTCAG-3' (SEQ ID NO:29);

p53 (300-393):

5'-ATGAATCCCCCAGGGAGCACT-3' (SEQ ID NO:30) and

5'-TGCGGCCGCTCAGTCTGAGTCAG-3' (SEQ ID NO:31).

The PCR products were cloned into the following restriction sites of the pGEX-5X1 GST vector: wt *p53*

EcoRI/NotI; *p53* (1-43) *EcoRI/NotI*; *p53* (1-200)

EcoRI/XhoI; *p53* (100-300) *EcoRI/XhoI*; *p53* (201-393)

EcoRI/NotI; *p53* (300-393) *EcoRI/NotI*. To make the

GST-BRCA1 (91-500) fusion protein, the *BRCA1* (91-500)

product was generated by PCR from the human *BRCA1* cDNA

with primers 5'-AGAATTCTGTGCTTTTCAGCTT-3' (SEQ ID NO:32)

and 5'-TCTCGAGTCAATTTGTGAGGGGACG-3' (SEQ ID NO:33).

After digestion with *EcoRI* and *XhoI*, the fragment was subcloned into pGEX-5X1. The cloned DNA sequence of all of the above vectors was verified.

Transfections and Luciferase Assays. SW480 and MEF cells were transfected using lipofectin and lipofectamine (GIBCO) respectively as described previously (El-Deiry et al., 1993, *supra*). SAOS2 cells
5 were transfected using the calcium phosphate precipitation described in Example 1. Luciferase assays were performed as described (Zeng et al., 1997, *supra*).

GST-fusion Protein Purification. Various pGEX-5X1 fusion constructs were transformed into DH5 α *E. coli* (GIBCO/BRL). The GST-p53 or GST-BRCA1 (91-500)
10 proteins were produced following incubation of the bacteria with 0.1-1 mM IPTG at 30°C for 2-3 hours. The cell pellet was resuspended in PBS and sonicated. The soluble GST-fusion proteins were immobilized onto
15 Glutathione Sepharose 4B beads (Pharmacia).

In vitro Interaction Between BRCA1 and p53. *In vitro* translations were carried out in the presence of 40 μ Ci [³⁵S]-methionine (NEN) using TNT7 coupled reticulocyte lysate system (Promega). For *in vitro* binding assays,
20 five to 20 μ l of *in vitro* translated ³⁵S-labeled proteins or cell lysate from 1x10⁷ cells was incubated with glutathione-sepharose beads containing GST-fusion proteins from 10 ml bacterial cultures. After washing, the bound proteins were eluted with 30 μ l SDS sample
25 buffer and resolved by 7.5% SDS-PAGE followed by fluorography. In experiments where unlabeled cell extracts were used, after SDS/PAGE separation, the proteins were transferred to Hybond™P membrane (Amersham). Immunoblotting was carried out using
30 anti-BRCA1 monoclonal antibody (SG11, 1:100 dilution; Oncogene Science) and ECL™ (Amersham).

TUNEL Assay. SW480 cells (3x10⁴/ well) were transfected with mammalian expression vectors (1 μ g) in

8-well chamber slides. The cells were fixed with neutral buffered formalin after 40 hrs and analyzed for fluorescence emission by TUNEL assay using the Apoptag Plus In Situ apoptosis detection kit-fluorescein (ONCOR) as previously described (Wu et al., 1997, Nat. Genetics 17:141-143). Briefly, fixed cells were incubated with digoxigenin-labeled nucleotides and TdT enzyme followed by washing with PBS and incubation with an anti-digoxigenin-FITC conjugate. In each experiment, the % of TdT(+) cells was determined by counting at least 300 cells from 5-10 random transfected fields.

Immunoprecipitations and Immunoblotting of BRCA1 and p53. Immunoprecipitation was carried out using an immunoprecipitation kit (Boehringer Mannheim). About 1×10^7 cells were lysed in Wash Buffer 1 (50 mM Tris, pH 7.5, 150 mM NaCl, 1% NP40, 0.5% sodium deoxycholate plus protease inhibitor cocktail). After preclearing with protein G-agarose beads, lysates were mixed with antibodies (1 μ g/ml) and protein G-agarose. The antibodies used are anti-BRCA1, (C20; Santa Cruz) and (SG11; Oncogene Science), anti-p16 (Ab1; Oncogene Science), anti-MDM-2 (Ab1; Oncogene Science) and anti-p53 (Ab-6; Oncogene Science). The beads were washed three times with Wash Buffer 1, one time with Wash Buffer 2 (50 mM Tris, pH 7.5, 500 mM NaCl, 0.1% NP40, 0.05% sodium deoxycholate), and one time with Wash Buffer 3 (50 mM Tris, pH 7.5, 0.1% NP40, 0.05% sodium deoxycholate). Samples were solubilized in SDS-sample buffer by boiling for 5 minutes and separated by 7.5 % SDS-PAGE. After transferring, the membranes were incubated with anti-p53 antibody conjugated to HRP (1:200 dilution of p53DO1; Santa Cruz) or anti-BRCA1 antibody (1:200 dilution of SG11; Oncogene Science). A 1:300 dilution of anti-mouse

HRP (Amersham) was used to detect BRCA1.

B. Results

BRCA1 stimulates p53-dependent transcription.

Because both p53 and BRCA1 can transactivate the p21
5 promoter, we investigated the effects of BRCA1 on
p53-dependent transcription. BRCA1 either alone or in
combination with p53 was cotransfected into SW480 colon
carcinoma cells together with a synthetic p53-specific
reporter (PG13-LUC) (El-Deiry et al., 1993, *supra*). The
10 expression of BRCA1 in SW480 (endogenous mutant p53) had
no effect on p53-specific reporter expression in the
absence of exogenous wild-type p53 (Fig. 5A). As
previously reported, transfection of p53 in SW480 cells
led to stimulation of the PG13-luciferase reporter
15 (El-Deiry et al., 1993, *supra*). Unexpectedly, a dramatic
increase in luciferase activity was observed following
cotransfection of SW480 cells by BRCA1 and wt p53
expression vectors (Fig. 5A). We confirmed the effect of
BRCA1 on p53-dependent transcription in MEF p53^{-/-} (Fig.
20 5B), wild-type p53-expressing HCT116 (Fig. 5C), wild-type
p53-expressing MCF7 breast cancer cells (Fig. 5D) and
SAOS-2 (p53-null Osteosarcoma) cells (data not shown).
The coactivation of p53-dependent transcription by BRCA1
was dose-dependent with respect to BRCA1 and required the
25 presence of wild-type p53 (Fig. 5E).

The transcriptional activity of p53 leads to
increased expression of p21 and bax, putative effectors
of p53-mediated growth arrest and apoptosis respectively
(Levine, 1997). BRCA1 stimulated p53-dependent
30 transcription by six- to eight-fold from the p21 and bax
promoters, respectively (Fig. 5F,G). As expected,
expression of BRCA1 alone activated the p21 promoter (see
Example 1). However, BRCA1 did not activate the bax

promoter in the absence of exogenous wt p53 (Fig. 5G, lane 2). Because of the increased expression of endogenous p53 target genes in wt p53-expressing cells as compared to cells which lack wt p53 (Wu et al., 1997, *supra*; El-Deiry et al., 1994, *supra*; Blasgosklonny et al., 1997, Intl. J. Oncology 11:1165-1170) and because of the lack of availability of BRCA1-null cell lines, it has been difficult to demonstrate that BRCA1 co-activates expression of endogenous p53 targets. It remains unclear whether endogenous p53-dependent p21 expression requires BRCA1. In transfection experiments it is possible to limit the amount of p53, a potent transcription factor, to demonstrate the role of BRCA1 as a coactivator. Because numerous studies have previously correlated the expression of endogenous genes with these reporters, we believe that the coactivation of p53-dependent transcription provides important insight into how BRCA1 may regulate the p53-dependent growth inhibitory pathway.

BRCA1 interacts with p53. BRCA1 may regulate p53-mediated transcription either directly through protein-protein interaction or indirectly through association with other proteins whose function may stimulate p53-mediated transcription. To determine whether BRCA1 binds to p53 *in vitro*, we used either glutathione S-transferase (GST) or GST-p53 fusion protein bound to Glutathione sepharose beads to precipitate either *in vitro* translated or cellular BRCA1 proteins. Approximately 20% of input ³⁵S-labeled BRCA1 bound to GST-p53 protein but no BRCA1 bound to GST alone. ³⁵S-labeled Luciferase protein, a negative control, did not bind to either GST or GST-p53.

In a similar experiment, COS-7 cell extracts were mixed with beads containing GST or GST-p53 and the

bound proteins were analyzed by BRCA1 immunoblotting. BRCA1 specifically associated with GST-p53 but not GST. To investigate the interaction between BRCA1 and p53 proteins *in vivo*, immunoprecipitations followed by western blots were performed. In these experiments, an anti-p53 monoclonal antibody directly conjugated to horseradish peroxidase was used to detect p53 directly. The endogenous p53 of COS-7, HeLa, U2OS and SW480 cells co-immunoprecipitated with BRCA1. Although p53 protein is well known to be targeted for degradation by HPV-E6 protein in HeLa cells, it was readily detectable in our co-immunoprecipitation assay. Interaction of p53 with CBP has also been demonstrated in HeLa cells (Gu et al., 1997, Nature 387:819-822). We have consistently observed that endogenous BRCA1 could be specifically recovered in association with the exogenous GST-p53 fusion protein linked to agarose beads.

We have found no evidence for a BRCA1-dependent supershift of p53-DNA binding site electrophoretic mobility using two different anti-human BRCA1 antibodies. It is possible that the interaction between BRCA1 and p53 may not be stable enough for gel shift experiments, that BRCA1 is not required for the DNA binding by p53, or that BRCA1 may be present but concealed by other proteins. Although BRCA1 and p53 appear to associate both *in vitro* and *in vivo*, we cannot exclude the possibility that the interaction may be indirect.

A novel N-terminal domain of BRCA1 exon 11 interacts with p53 *in vitro*. To identify the region of BRCA1 that interacts with p53, a series of BRCA1 deletion mutants (Fig. 6) translated *in vitro* in the presence of ³⁵S-methionine were individually mixed with GST-p53-bound affinity resin. The region of BRCA1 extending from aa

residues 224 to 500 was found to be sufficient for interaction with p53. This region corresponds to the N-terminal region of exon 11 of BRCA1. To further confirm the minimal BRCA10-interacting region, either GST or GST-BRCA1 fusion protein containing residues 74 to 500 of BRCA1, was mixed with *in vitro* synthesized full length p53 protein. p53 protein specifically bound to the GST-BRCA1, whereas luciferase protein, did not bind to either GST or GST-BRCA1. This conserved domain (Szabo et al., 1996, Hum. Mol. Genet. 5:1289-1298) has not been noted previously to play a role in BRCA1 function. It is notable that this region is missing from several alternatively spliced forms of BRCA1 (Thakur et al., 1997, *supra*).

BRCA1 binds to the C-terminal domain of p53.
In order to identify the region of p53 (Ko and Prives, 1996, Genes Dev. 10:1054-1072) which binds to BRCA1, a series of p53 deletion mutants (Fig. 7) were generated as GST-fusion proteins and their ability to interact with *in vitro* translated full-length BRCA1 or exon 11 of BRCA1 protein was studied. The region of p53 between aa residues 300-393 was found to be sufficient for its interaction with BRCA1. These results suggest that the physical interaction between p53 and BRCA1 may underlie the BRCA1-dependent stimulation of p53-mediated transcription.

BRCA1 mutants are deficient in stimulation of p53-dependent transcription. To further explore the biological relevance of the stimulation of p53-dependent transcription by BRCA1, we studied the effect of various synthetic and tumor-derived mutant BRCA1 proteins on p53-dependent transcriptional activity. Mutants of BRCA1 lacking the C-terminus transactivation domain or both the

C-terminus transactivation domain and the Rad51-interacting domain were poor augmenters of p53-mediated transcriptional activity (Fig. 8A). Similarly, four tumor-derived transactivation-deficient *BRCA1* mutants were defective in stimulation of p53-mediated transcription (Fig. 8A). These results suggest that an intact transactivation domain within *BRCA1* may be required for *BRCA1* to increase p53-dependent transcriptional activity.

10 *BRCA1* regulates p21^{WAF1/CIP1} through a p53-dependent as well as a p53-independent mechanism. p53 had no effect on *BRCA1*-dependent activation of the p21-promoter in the absence of p53 DNA-binding sites (Fig. 8B), suggesting that the effect of *BRCA1* on p53 (Fig. 5) is specific to p53-up-regulated genes. Thus it is clear that *BRCA1* can activate p21 by a p53-independent pathway that maps to a region distinct from the p53 DNA binding site and can coactivate p53-dependent gene expression in general only in the presence of p53 binding sites (Figs. 1, 4).

20 Dominant negative inhibition of p53-dependent transcriptional activity by a *BRCA1* truncation mutant retaining the p53-interacting domain. The *BRCA1* mutant Δ500-1863 (Example 1), which lacks both the Rad51-interacting and C-terminal transactivation domains, was used to test whether truncating *BRCA1* mutants retaining the newly-identified p53-interacting domain may act as dominant negative inhibitors of p53-mediated transcription. We predicted that this truncated *BRCA1* protein might inhibit p53-mediated transcriptional activity by competing with endogenous *BRCA1* for p53. Co-transfection of truncated *BRCA1* (Δ500-1863) with p53 inhibited p53-mediated transcription in a dose-dependent

manner (Fig. 8C). The apparent dominant negative inhibition of p53 was overcome by increasing amounts of exogenous wt BRCA1 (Fig. 8C). It may be that the truncated BRCA1 protein, which does not have a nuclear localization signal, may associate with p53 prior to its entry into the nucleus.

Cooperation between BRCA1 and p53 in apoptosis induction. p53-mediated apoptosis plays a major role in tumor suppression. One mechanism by which p53 mediates apoptosis is by its ability to activate transcription (Haupt et al., 1995, Genes Dev. 9:2170-2183; Chen et al., 1996, Genes Dev. 10:2438-2451). In order to determine if BRCA1 has any effect on p53-mediated apoptosis through its ability to stimulate p53-mediated transcription, p53 was co-transfected with either BRCA1 or its vector into SW480 cells. Apoptotic cells were identified by TUNEL assay and the apoptotic cells (%) was quantified at 40 hrs following transfection (Table 2). Vector-alone transfected cells showed a low level of apoptosis. Transfection of p53 or BRCA1 alone induced apoptosis up to 10-fold and co-transfection of BRCA1 with p53 increased further the apoptotic cells (Table 2) suggesting p53 and BRCA1 may cooperate to induce apoptosis. SW480 cells express low levels of endogenous BRCA1 (Example 1), and so it is possible that the observed levels of apoptosis following p53 transfection alone may already involve cooperation with BRCA1. The modest increase in apoptosis due to exogenous BRCA1 should also be considered in the context that upper limit for (%) apoptosis (range of 20-50%) in these experiments is the transfection efficiency. The results suggest a scenario wherein BRCA1 may enhance the induction of apoptosis by p53 possibly through

stimulation of p53's transcriptional activity.

Table 2.

5	Apoptotic Cells (%)			
	Plasmids* Used	Expt. #1	Expt. #2	Expt #3
10	pCEP4+pCR3	2.3	2.1	1.9
	pCEP4+BRCA1	9.2	7.0	16.6
	p53+pCR3	3.4	20.7	11.9
15	p53+BRCA1	11.8	29.1	18.0

*SW480 cells were transfected with p53 (0.5 μ g) or its vector pCEP4 and BRCA1 (0.5 μ g) or its vector pCR3 (0.5 μ g) in different combinations. Apoptotic cells were detected by TUNEL assay as described in Materials and Methods. Apoptotic cells show strong fluorescence staining of fragmented chromatin.

C. Discussion

Both p53 and BRCA1 are physically altered by the cellular response to DNA damage, p53 by stabilization and BRCA1 by hyperphosphorylation (Kastan et al., 1991, Cancer Res. 51:6304-6311; Scully et al., 1997, Cell 90:425-435). Both proteins can activate expression of the cell cycle inhibitor p21 (El-Deiry et al., 1993, supra; Example 1). Because both BRCA1 and p53 are implicated in regulation of gene transcription, control of cell growth, and response to DNA damage, we sought to investigate the possibility of a functional cross-talk between these two proteins. The results suggest that BRCA1 and p53 can physically associate, both *in vitro* and *in vivo*, and function in a common pathway of tumor suppression. The ability of BRCA1 to biochemically modulate p53 function suggests that this may be a fundamental role of BRCA1 in tumor suppression. It is

noteworthy that inherited defects of either p53 or BRCA1 can lead to the development of breast cancer in humans (Malkin et al., 1990, Science 250:1233-1238; Miki et al., 1994, Science 266:66-71).

5 The results reported in this example reveal that BRCA1 is a potent activator of p53-dependent transcription by demonstrating that BRCA1 potentiates p53-activated transcription of the *p21* and the *bax* promoters. The role of p53 as a transcription factor is
10 well established (Vogelstein and Kinzler, 1992, Cell 70:523-526) and appears to be important for its ability to arrest cell cycle progression as well as to induce apoptosis (Levine, 1997, Cell 88:323-331). BRCA1 has been postulated to function as a coactivator of
15 transcription, based on its localization as a component of RNA polymerase II holoenzyme and transactivation function (Example 1 and Chapman and Verma, 1996, Nature 382:678-679; Monteiro et al., 1996, PNAS 93:13595-13599; Scully et al., 1997, PNAS 94:5605-5610). The observation
20 that BRCA1 is a transcriptional coactivator of p53, including strong up-regulation of expression of the *bax* gene, provides a novel mechanism for apoptosis induction and tumor suppression by BRCA1. BRCA1 serves a co-stimulatory role in p53-mediated transcription from
25 *p21*, suggesting BRCA1 is involved in both p53-dependent transcription and p53-independent transcription. This finding is consistent with previous data demonstrating that *p21* can be activated by other p53 independent pathways (El Deiry et al., 1998, Curr. Top. Micro.
30 Immunol. 227:121-137).

During gene expression, a complex of activators, coactivators, and basal factors are necessary to stabilize the interactants and provide transcriptional

specificity. BRCA1 may link a variety of different DNA binding factors such as p53 to the components of the basal transcription machinery. Here we show that p53 and BRCA1 form a specific protein complex *in vitro* and *in vivo*. An N-terminal segment of BRCA1 exon 11 extending from residues 224 to 500 is sufficient for its interaction with p53. Interestingly, exon 11 is deleted in several naturally occurring isoforms of BRCA1 (Thakur et al, 1997, *supra*), leading to speculation that the inability of the BRCA1Δ exon11 mutant to bind to p53 may be a regulatory mechanism for BRCA1. Our finding also suggests that BRCA1 binding to p53 may not be mediated by RAD51, RNA polymerase II holoenzyme or BARD1 since they bind to different regions on BRCA1 (Scully et al, 1997, *supra*; Scully et al, 1997, Cell 88:265-275; Wu et al, 1996, Nature Genet. 14:430-440). Post-translational modifications of the C-terminus of p53 have been shown to play an important role in controlling p53-specific DNA binding (Levine, 1997, Cell 88:323-331). Phosphorylation, acetylation, antibody binding, or deletion of this region can convert p53 from an inert to an active form for DNA binding (Ko and Prives, 1997, *supra*; Gu and Roeder, 1997, Cell 90:595-606). Therefore allosteric regulation of p53 by distinct cellular signaling pathways modulates the conversion between latent and activated forms of p53. Our results show BRCA1 binds to the C-terminus of p53. BRCA1 may activate p53 by altering the conformation of p53 through direct physical binding and/or modification of the C-terminal p53. In addition, p53 may be bridged to the basal transcriptional machinery by BRCA1.

Mutation in the C-terminus of BRCA1, which contains its transactivation domain (Chapman and Verma,

1996, *supra*; Monteina et al., 199, *supra*), occurs in hereditary breast cancer (Couch and Weber, 1994, Human Mutation 8:8-18). Such mutants of BRCA1 are defective in the ability to activate $p21^{RAF1/CIP1}$ (Example 1). The results reported in this example indicate that the tumor derived BRCA1 mutants carrying point mutations in the C-terminus are also defective in stimulation of p53-dependent transcription. Because BRCA1 interaction with p53 maps to the N-terminal region of BRCA1, we predicted that truncated-BRCA1 proteins, the most common forms found in hereditary breast and ovarian cancer, may compete for sites on p53 and perhaps inhibit p53 function. We found that truncated BRCA1 is a potent dominant negative inhibitor of p53-dependent transcription, and that this could be reversed by excess wild-type BRCA1. These results would predict a defective p53 transcriptional activity in certain cancer cells.

Although p53 and BRCA1 appear to physically interact directly *in vitro*, we found no evidence for stimulation of BRCA1-dependent transactivation of $p21$ by exogenous p53 in the absence of p53-DNA binding sites. BRCA1 could, however, greatly stimulate p53-dependent activation of the *bax* promoter, in the absence of p53-independent activation of this promoter by BRCA1. These observations argue that, *in vivo*, certain DNA regulatory elements serve to recruit p53 to certain genes. In such situations the presence of p53 protein favors the chance for local association with BRCA1 and cooperative activation is observed. In our system using either the $p21$ or *bax* promoter, the combination of p53 and BRCA1 leads to more than an additive increase in gene expression, but it is not clear whether BRCA1 is required for p53-dependent transcription, because of the lack of

availability of BRCA1-null cell lines. The
BRCA1-knockout embryos appear to express p21 (Hakem et
al., 1996, Cell 85:1009-1023) and the current model is
that this may be due to genomic instability and
5 subsequent p53 activation (Brugarolas and Jacks, 1997,
Nature Med. 7:721-722). The dominant negative action of
truncated BRCA1 on p53-dependent transcription hints at
the possibility that BRCA1 may be required for
p53-dependent transcription. Our results argue that in
10 somatic cells, BRCA1 can activate p53 function leading to
tumor suppression.

The induction of apoptosis by p53 is critical
for its tumor suppressor function (Symonds et al., 1994,
Cell 78:703-711). The results presented in Table 1
15 suggest that BRCA1 cooperates with p53 in inducing
apoptosis. Altogether, our data suggest that BRCA1 acts
as a coactivator of p53-driven gene expression and may
play an important role in cell growth arrest, apoptosis,
and/or DNA damage repair processes. It has not been
20 determined if BRCA1 is required for transcription-
independent p53 action. The possibility that the complex
of BRCA1 and p53 may directly participate in DNA damage
recognition and/or repair could not be excluded. In
summary, we have identified a physical and functional
25 association between the tumor suppressors p53 and BRCA1,
linking their biochemical effects to a common pathway of
tumor suppression.

~~The present invention is not limited to the~~
30 ~~embodiments described and exemplified above, but is~~
~~capable of variation and modification without departure~~
~~from the scope of the appended claims.~~

We claim:

1. An isolated nucleic acid molecule comprising a region of an upstream promoter of a *p21* gene, that contains an element required for p53-independent BRCA1-mediated transactivation of the gene.
5
2. The nucleic acid molecule of claim 1, wherein the region is located on a human *p21* gene between the transcription start site and 143 nucleotides 5' of the transcription start site of the gene.
10
3. The nucleic acid molecule of claim 2, wherein the region comprises nucleotides 4441-4584 of SEQ ID NO:3.
15
4. An isolated nucleic acid molecule comprising a region of an upstream promoter of a *p21* gene, the removal of which is associated with prevention of p53-independent transactivation of the gene by BRCA1.
20
5. The nucleic acid molecule of claim 1, wherein the region is located on a human *p21* gene between 93 and 143 nucleotides 5' of the transcription start site of the gene.
25
6. The nucleic acid molecule of claim 2, wherein the region comprises nucleotides 4441-4491 of SEQ ID NO:3.
7. An isolated polypeptide comprising a region of a BRCA1 protein that physically interacts with a p53 protein.
30

8. The polypeptide of claim 7, wherein the region resides within amino acid residues 224-500 of a human BRCA1 protein.

5 9. The polypeptide of claim 8, wherein the human BRCA1 protein has the amino acid sequence of SEQ ID NO:1.

10 10. Antibodies immunologically specific for the polypeptide of claim 7.

11. An isolated nucleic acid encoding the polypeptide of claim 7.

15 12. A vector comprising the nucleic acid of claim 11.

20 13. An isolated polypeptide comprising a region of a p53 protein that physically interacts with a BRCA1 protein.

25 14. The polypeptide of claim 13, wherein the region resides within amino acid residues 300-395 of a human p53 protein.

15. The polypeptide of claim 4, wherein the human BRCA1 protein has the amino acid sequence of SEQ ID NO:2.

30 16. Antibodies immunologically specific for the polypeptide of claim 13.

17. An isolated nucleic acid molecule encoding the polypeptide of claim 13.

18. A vector comprising the nucleic acid molecule of claim 17.

19. An isolated protein-protein complex comprising a p53 protein and a BRCA1 protein.

20. The complex of claim 19, comprising human p53 and human BRCA1.

21. A method for determining if a test compound enhances or interferes with formation of a complex comprising BRCA1 and p53, which comprises:

a) providing an assay test mixture comprising the BRCA1, the p53 and the test compound;

b) providing an equivalent assay control mixture comprising the BRCA1 and the p53, without the test compound;

c) subjecting the test mixture and the control mixture to conditions known to enable formation of the BRCA1-p53 complex in the absence of the test compound;

d) measuring the amount of BRCA1-p53 complexes formed, if any, in the test mixture and the control mixture; and

e) determining if the amount of BRCA1-p53 complex formed in the test mixture is different from the amount of the complex formed in the control mixture, an increase being indicative of the ability of the test compound to enhance formation of the BRCA1-p53 complex and a decrease being indicative of the ability of the test compound to interfere with formation of the BRCA1-p53 complex.

22. The method of claim 21, wherein the assay mixtures are cell-free assay mixtures and the BRCA1 and p53 proteins are provided as isolated proteins.

5

23. The method of claim 21, wherein the assay mixtures comprise cells, and the BRCA1 and p53 proteins are provided by expression of nucleic acid molecules within the cells.

10

24. A method for determining if a test compound enhances or interferes with p53-independent BRCA1-mediated transactivation of a *p21* gene, which comprises:

15

a) providing a cell line deficient in production of endogenous functional BRCA1 and p53, comprising at least one heterologous vector expressing BRCA1, said cell further comprising a reporter gene operably linked to a *p21* promoter containing a region required for the p53-independent BRCA1-mediated transcription, wherein the reporter gene expresses a detectable gene product as a result BRCA1-mediated transactivation of the *p21* promoter;

20

b) preparing a culture of the cells;

25

c) incorporating the test compound into the cell culture under conditions permitting expression of the heterologous vectors and the reporter gene, and BRCA1-mediated transactivation of the *p21* promoter; and

d) measuring expression of the reporter

~~30~~ gene by detecting the presence or amount, if any, of the detectable gene product and comparing the expression to an equivalent cell culture in which the test compound was not incorporated, an increase in expression of the reporter gene in the culture containing the test compound

being indicative that the test compound enhances p53-independent BRCA1-mediated transactivation of the p21 gene, a decrease in expression of the reporter gene being indicative that the test compound interferes with p53-independent BRCA1-mediated transactivation of the p21 gene.

25. The method of claim 24, wherein the cells lack endogenous functional proteins selected from the group consisting of BRCA1 and p53.

26. A method for determining if a test compound enhances or interferes with tumor suppression resulting from p53-independent BRCA1-mediated transactivation of a p21 gene, which comprises:

- a) providing a tumor cell line comprising at least one heterologous vector expressing BRCA1;
- b) preparing a culture of the cells;
- c) incorporating the test compound into the cell culture under conditions permitting expression of the heterologous vector and BRCA1-mediated transactivation of an endogenous p21 gene; and
- d) determining proliferation of the cell culture and comparing it to proliferation of an equivalent cell culture in which the test compound was not incorporated, an increase in the proliferation of the culture containing the test compound being indicative that the test compound interferes with tumor suppression, a decrease in the proliferation of the culture containing the test compound being indicative that the test compound enhances tumor suppression.

27. The method of claim 26, wherein the cells lack endogenous functional proteins selected from the group consisting of BRCA1 and p53.

28. The method of claim 26, wherein the proliferation rate is measured by measuring DNA synthesis in the cultured cells.

5

29. A method for determining if a test compound enhances or interferes with p53-dependent BRCA1-mediated transactivation of a gene, which comprises:

a) providing a cell line comprising at
10 least one heterologous vector expressing BRCA1 and p53, said cell further comprising a reporter gene operably linked to a promoter of a gene comprising sequences enabling p53-dependent BRCA1-mediated transactivation of the gene, wherein the reporter gene expresses a
15 detectable gene product as a result BRCA1-mediated transactivation of the p21 promoter;

b) preparing a culture of the cells;

c) incorporating the test compound into
the cell culture under conditions permitting expression
20 of the heterologous vectors, reporter gene and p53-dependent BRCA1-mediated transactivation of the promoter; and

d) measuring expression of the reporter gene by detecting the presence or amount, if any, of the
25 detectable gene product and comparing the expression to an equivalent cell culture in which the test compound was not incorporated, an increase in expression of the reporter gene in the culture containing the test compound being indicative that the test compound enhances p53-
30 dependent BRCA1-mediated transactivation of the gene, a decrease in expression of the reporter gene being indicative that the test compound interferes with p53-dependent BRCA1-mediated transactivation of the gene.

30. The method of claim 29, wherein the cells lack endogenous functional proteins selected from the group consisting of BRCA1 and p53.

5 31. A method for determining if a test compound enhances or interferes with tumor suppression resulting from p53-dependent BRCA1-mediated transactivation of a tumor suppressor gene, which comprises:

10 a) providing a tumor cell line comprising at least one heterologous vector expressing BRCA1 and p53;

 b) preparing a culture of the cells;

 c) incorporating the test compound into
15 the cell culture under conditions permitting expression of the heterologous vectors and p53-dependent BRCA1-mediated transactivation of the gene; and

 d) determining proliferation of the cell culture and comparing it to proliferation of an
20 equivalent cell culture in which the test compound was not incorporated, an increase in the proliferation of the culture containing the test compound being indicative that the test compound interferes with tumor suppression, a decrease in the proliferation of the culture containing
25 the test compound being indicative that the test compound enhances tumor suppression.

32. The method of claim 31, wherein the cells lack endogenous functional proteins selected from the
30 group consisting of BRCA1 and p53.

33. The method of claim 31, wherein the gene is selected from the group consisting of *bax* and *p21*.

34. The method of claim 31, wherein the proliferation is determined by measuring DNA synthesis in the cultured cells.

5 35. The method of claim 31, wherein the proliferation is determined by measuring apoptosis in the cultured cells.

10 36. A kit for performing the method of claim 21, comprising a container containing at least one biological molecule selected from the group consisting of BRCA1 protein, p53 protein, an expressible nucleic acid molecule encoding BRCA1, an expressible nucleic acid molecule encoding p53, and a reporter gene.

15 37. A kit for performing the method of claim 24, comprising a container containing at least one biological molecule selected from the group consisting of an expressible vector encoding BRCA1, and the reporter gene.

20 38. The kit of claim 37, which further comprises the tumor cell line.

25 39. A kit for performing the method of claim 26, which comprises a container containing an expressible vector encoding BRCA1.

30 ~~40. The kit of claim 39, which further~~
comprises the tumor cell line.

41. A kit for performing the method of claim 29, comprising a container containing at least one

biological molecule selected from the group consisting of an expressible vector encoding BRCA1, an expressible vector encoding p53, and the reporter gene.

5 42. The kit of claim 41, which further comprises the tumor cell line.

 43. A kit for performing the method of claim 31, which comprises a container containing an expressible
10 vector encoding BRCA1 and an expressible vector encoding p53.

 44. The kit of claim 43, which further comprises the tumor cell line.

15 45. A method for diagnosing a subject's predisposition for developing a tumor, comprising detecting defects in the subject's *BRCA1* genes, *BRCA1* proteins or *p21* promoters that result in reduction or
20 prevention of p53-independent *BRCA1*-mediated transactivation of the subject's *p21* genes.

 46. A method for diagnosing a subject's predisposition for developing a tumor, comprising
25 detecting defects in the subject's *BRCA1* genes, *BRCA1* proteins, *p53* genes, *p53* proteins, or promoters of tumor suppressor genes co-activated by *BRCA1* and *p53*, that result in reduction or prevention of p53-dependent *BRCA1*-mediated transactivation of the tumor suppressor genes.


30

 47. A method for enhancing tumor suppression which comprises increasing p53-dependent or -independent *BRCA1*-mediated transactivation of at least one tumor

suppressor gene in tumor cells.

48. A method for increasing proliferation of cells, which comprises decreasing p53-dependent or -
5 independent BRCA1-mediated transactivation of at least one tumor suppressor gene in the cells.

Fig. 1



2/10

Fig. 2A




Fig. 2B





Fig. 2C



3/10

Fig. 3A




Fig. 3B






Fig. 3C



4/10

Fig. 4



5/10

Fig. 5A




Fig. 5B




Fig. 5C





Fig. 5D



6/10

Fig. 5E




Fig. 5F




Fig. 5G




Fig. 6





Fig. 7



9/10

Fig. 8A





Fig. 8B



10/10

Fig. 8C





Fig. 8D



SEQUENCE LISTING

5 <110> The Trustees of the University of Pennsylvania
 El-Deiry, W.S.
 Weber, B.L.

10 <120> Compositions and Methods for Controlling
 BCRA1-mediated p53-Dependent and -Independent Regulation of
 Transcription

<130> Penn K-1825

15 <150> 60/080,146
 <151> 1998-03-31

<160> 33

20 <170> FastSEQ for Windows Version 3.0

<210> 1
 <211> 1863
 <212> PRT
 <213> Homo sapiens

25 <220>
 <221> BINDING
 <222> (224)...(500)
 <223> /note="Binding site of p53 protein"

30 <300>
 <301> Smith, T.M. et al.,
 <302> Complete genomic seq. and analysis of 117 kb of hu
 <303> Genome Research

35 <304> 6
 <305> 11
 <306> 1029-1049
 <307> 1996-01-01
 <308> Genbank L78833

40 <309> 1996-12-04

<400> 1
 Met Asp Leu Ser Ala Leu Arg Val Glu Glu Val Gln Asn Val Ile Asn
 1 5 10 15
 45 Ala Met Gln Lys Ile Leu Glu Cys Pro Ile Cys Leu Glu Leu Ile Lys
 20 25 30
 Glu Pro Val Ser Thr Lys Cys Asp His Ile Phe Cys Lys Phe Cys Met
 35 40 45
 Leu Lys Leu Leu Asn Gln Lys Lys Gly Pro Ser Gln Cys Pro Leu Cys
 50 50 55 60
 Lys Asn Asp Ile Thr Lys Arg Ser Leu Gln Glu Ser Thr Arg Phe Ser
 65 70 75 80
 Gln Leu Val Glu Glu Leu Leu Lys Ile Ile Cys Ala Phe Gln Leu Asp
 85 90 95
 55 Thr Gly Leu Glu Tyr Ala Asn Ser Tyr Asn Phe Ala Lys Lys Glu Asn
 100 105 110
 Asn Ser Pro Glu His Leu Lys Asp Glu Val Ser Ile Ile Gln Ser Met
 115 120 125
 Gly Tyr Arg Asn Arg Ala Lys Arg Leu Leu Gln Ser Glu Pro Glu Asn

	130		135		140	
	Pro Ser Leu Gln Glu Thr Ser Leu Ser Val Gln Leu Ser Asn Leu Gly					
	145		150		155	160
5	Thr Val Arg Thr Leu Arg Thr Lys Gln Arg Ile Gln Pro Gln Lys Thr					
		165		170		175
	Ser Val Tyr Ile Glu Leu Gly Ser Asp Ser Ser Glu Asp Thr Val Asn					
		180		185		190
	Lys Ala Thr Tyr Cys Ser Val Gly Asp Gln Glu Leu Leu Gln Ile Thr					
		195		200		205
10	Pro Gln Gly Thr Arg Asp Glu Ile Ser Leu Asp Ser Ala Lys Lys Ala					
		210		215		220
	Ala Cys Glu Phe Ser Glu Thr Asp Val Thr Asn Thr Glu His His Gln					
	225		230		235	240
	Pro Ser Asn Asn Asp Leu Asn Thr Thr Glu Lys Arg Ala Ala Glu Arg					
15		245		250		255
	His Pro Glu Lys Tyr Gln Gly Ser Ser Val Ser Asn Leu His Val Glu					
		260		265		270
	Pro Cys Gly Thr Asn Thr His Ala Ser Ser Leu Gln His Glu Asn Ser					
		275		280		285
20	Ser Leu Leu Leu Thr Lys Asp Arg Met Asn Val Glu Lys Ala Glu Phe					
		290		295		300
	Cys Asn Lys Ser Lys Gln Pro Gly Leu Ala Arg Ser Gln His Asn Arg					
	305		310		315	320
	Trp Ala Gly Ser Lys Glu Thr Cys Asn Asp Arg Arg Thr Pro Ser Thr					
25		325		330		335
	Glu Lys Lys Val Asp Leu Asn Ala Asp Pro Leu Cys Glu Arg Lys Glu					
		340		345		350
	Trp Asn Lys Gln Lys Leu Pro Cys Ser Glu Asn Pro Arg Asp Thr Glu					
		355		360		365
30	Asp Val Pro Trp Ile Thr Leu Asn Ser Ser Ile Gln Lys Val Asn Glu					
		370		375		380
	Trp Phe Ser Arg Ser Asp Glu Leu Leu Gly Ser Asp Asp Ser His Asp					
	385		390		395	400
	Gly Glu Ser Glu Ser Asn Ala Lys Val Ala Asp Val Leu Asp Val Leu					
35		405		410		415
	Asn Glu Val Asp Glu Tyr Ser Gly Ser Ser Glu Lys Ile Asp Leu Leu					
		420		425		430
	Ala Ser Asp Pro His Glu Ala Leu Ile Cys Lys Ser Glu Arg Val His					
		435		440		445
40	Ser Lys Ser Val Glu Ser Asn Ile Glu Asp Lys Ile Phe Gly Lys Thr					
		450		455		460
	Tyr Arg Lys Lys Ala Ser Leu Pro Asn Leu Ser His Val Thr Glu Asn					
	465		470		475	480
	Leu Ile Ile Gly Ala Phe Val Thr Glu Pro Gln Ile Ile Gln Glu Arg					
45		485		490		495
	Pro Leu Thr Asn Lys Leu Lys Arg Lys Arg Arg Pro Thr Ser Gly Leu					
		500		505		510
	His Pro Glu Asp Phe Ile Lys Lys Ala Asp Leu Ala Val Gln Lys Thr					
		515		520		525
50	Pro Glu Met Ile Asn Gln Gly Thr Asn Gln Thr Glu Gln Asn Gly Gln					
		530		535		540
	Val Met Asn Ile Thr Asn Ser Gly His Glu Asn Lys Thr Lys Gly Asp					
	545		550		555	560
55	Ser Ile Gln Asn Glu Lys Asn Pro Asn Pro Ile Glu Ser Leu Glu Lys					
		565		570		575
	Glu Ser Ala Phe Lys Thr Lys Ala Glu Pro Ile Ser Ser Ser Ile Ser					
		580		585		590
	Asn Met Glu Leu Glu Leu Asn Ile His Asn Ser Lys Ala Pro Lys Lys					
		595		600		605

	Asn	Arg	Leu	Arg	Arg	Lys	Ser	Ser	Thr	Arg	His	Ile	His	Ala	Leu	Glu	
	610						615					620					
	Leu	Val	Val	Ser	Arg	Asn	Leu	Ser	Pro	Pro	Asn	Cys	Thr	Glu	Leu	Gln	
	625					630					635					640	
5	Ile	Asp	Ser	Cys	Ser	Ser	Ser	Glu	Glu	Ile	Lys	Lys	Lys	Lys	Tyr	Asn	
					645					650					655		
	Gln	Met	Pro	Val	Arg	His	Ser	Arg	Asn	Leu	Gln	Leu	Met	Glu	Gly	Lys	
				660					665					670			
10	Glu	Pro	Ala	Thr	Gly	Ala	Lys	Lys	Ser	Asn	Lys	Pro	Asn	Glu	Gln	Thr	
			675					680					685				
	Ser	Lys	Arg	His	Asp	Ser	Asp	Thr	Phe	Pro	Glu	Leu	Lys	Leu	Thr	Asn	
	690					695					700						
	Ala	Pro	Gly	Ser	Phe	Thr	Lys	Cys	Ser	Asn	Thr	Ser	Glu	Leu	Lys	Glu	
	705				710					715						720	
15	Phe	Val	Asn	Pro	Ser	Leu	Pro	Arg	Glu	Glu	Lys	Glu	Glu	Lys	Leu	Glu	
					725				730						735		
	Thr	Val	Lys	Val	Ser	Asn	Asn	Ala	Glu	Asp	Pro	Lys	Asp	Leu	Met	Leu	
				740					745					750			
20	Ser	Gly	Glu	Arg	Val	Leu	Gln	Thr	Glu	Arg	Ser	Val	Glu	Ser	Ser	Ser	
		755					760						765				
	Ile	Ser	Leu	Val	Pro	Gly	Thr	Asp	Tyr	Gly	Thr	Gln	Glu	Ser	Ile	Ser	
	770					775					780						
	Leu	Leu	Glu	Val	Ser	Thr	Leu	Gly	Lys	Ala	Lys	Thr	Glu	Pro	Asn	Lys	
	785				790					795						800	
25	Cys	Val	Ser	Gln	Cys	Ala	Ala	Phe	Glu	Asn	Pro	Lys	Gly	Leu	Ile	His	
					805					810					815		
	Gly	Cys	Ser	Lys	Asp	Asn	Arg	Asn	Asp	Thr	Glu	Gly	Phe	Lys	Tyr	Pro	
				820					825					830			
30	Leu	Gly	His	Glu	Val	Asn	His	Ser	Arg	Glu	Thr	Ser	Ile	Glu	Met	Glu	
		835					840						845				
	Glu	Ser	Glu	Leu	Asp	Ala	Gln	Tyr	Leu	Gln	Asn	Thr	Phe	Lys	Val	Ser	
	850					855					860						
	Lys	Arg	Gln	Ser	Phe	Ala	Pro	Phe	Ser	Asn	Pro	Gly	Asn	Ala	Glu	Glu	
	865				870					875						880	
35	Glu	Cys	Ala	Thr	Phe	Ser	Ala	His	Ser	Gly	Ser	Leu	Lys	Lys	Gln	Ser	
					885					890					895		
	Pro	Lys	Val	Thr	Phe	Glu	Cys	Glu	Gln	Lys	Glu	Glu	Asn	Gln	Gly	Lys	
				900					905					910			
40	Asn	Glu	Ser	Asn	Ile	Lys	Pro	Val	Gln	Thr	Val	Asn	Ile	Thr	Ala	Gly	
		915					920						925				
	Phe	Pro	Val	Val	Gly	Gln	Lys	Asp	Lys	Pro	Val	Asp	Asn	Ala	Lys	Cys	
	930					935						940					
	Ser	Ile	Lys	Gly	Gly	Ser	Arg	Phe	Cys	Leu	Ser	Ser	Gln	Phe	Arg	Gly	
	945				950					955						960	
45	Asn	Glu	Thr	Gly	Leu	Ile	Thr	Pro	Asn	Lys	His	Gly	Leu	Leu	Gln	Asn	
					965					970					975		
	Pro	Tyr	Arg	Ile	Pro	Pro	Leu	Phe	Pro	Ile	Lys	Ser	Phe	Val	Lys	Thr	
				980					985					990			
50	Lys	Cys	Lys	Lys	Asn	Leu	Leu	Glu	Glu	Asn	Phe	Glu	Glu	His	Ser	Met	
			995				1000						1005				
	Ser	Pro	Glu	Arg	Glu	Met	Gly	Asn	Glu	Asn	Ile	Pro	Ser	Thr	Val	Ser	
	1010					1015					1020						
	Thr	Ile	Ser	Arg	Asn	Asn	Ile	Arg	Glu	Asn	Val	Phe	Lys	Glu	Ala	Ser	
	1025				1030						1035				1040		
55	Ser	Ser	Asn	Ile	Asn	Glu	Val	Gly	Ser	Ser	Thr	Asn	Glu	Val	Gly	Ser	
					1045					1050					1055		
	Ser	Ile	Asn	Glu	Ile	Gly	Ser	Ser	Asp	Glu	Asn	Ile	Gln	Ala	Glu	Leu	
				1060					1065					1070			
	Gly	Arg	Asn	Arg	Gly	Pro	Lys	Leu	Asn	Ala	Met	Leu	Arg	Leu	Gly	Val	

		1075		1080		1085	
		Leu Gln Pro Glu Val Tyr Lys Gln Ser Leu Pro Gly Ser Asn Cys Lys					
		1090		1095		1100	
5		His Pro Glu Ile Lys Lys Gln Glu Tyr Glu Glu Val Val Gln Thr Val					
		1105		1110		1115	1120
		Asn Thr Asp Phe Ser Pro Tyr Leu Ile Ser Asp Asn Leu Glu Gln Pro					
		1125		1130		1135	
		Met Gly Ser Ser His Ala Ser Gln Val Cys Ser Glu Thr Pro Asp Asp					
		1140		1145		1150	
10		Leu Leu Asp Asp Gly Glu Ile Lys Glu Asp Thr Ser Phe Ala Glu Asn					
		1155		1160		1165	
		Asp Ile Lys Glu Ser Ser Ala Val Phe Ser Lys Ser Val Gln Lys Gly					
		1170		1175		1180	
15		Glu Leu Ser Arg Ser Pro Ser Pro Phe Thr His Thr His Leu Ala Gln					
		1185		1190		1195	1200
		Gly Tyr Arg Arg Gly Ala Lys Lys Leu Glu Ser Ser Glu Glu Asn Leu					
		1205		1210		1215	
		Ser Ser Glu Asp Glu Glu Leu Pro Cys Phe Gln His Leu Leu Phe Gly					
		1220		1225		1230	
20		Lys Val Asn Asn Ile Pro Ser Gln Ser Thr Arg His Ser Thr Val Ala					
		1235		1240		1245	
		Thr Glu Cys Leu Ser Lys Asn Thr Glu Glu Asn Leu Leu Ser Leu Lys					
		1250		1255		1260	
25		Asn Ser Leu Asn Asp Cys Ser Asn Gln Val Ile Leu Ala Lys Ala Ser					
		1265		1270		1275	1280
		Gln Glu His His Leu Ser Glu Glu Thr Lys Cys Ser Ala Ser Leu Phe					
		1285		1290		1295	
		Ser Ser Gln Cys Ser Glu Leu Glu Asp Leu Thr Ala Asn Thr Asn Thr					
		1300		1305		1310	
30		Gln Asp Pro Phe Leu Ile Gly Ser Ser Lys Gln Met Arg His Gln Ser					
		1315		1320		1325	
		Glu Ser Gln Gly Val Gly Leu Ser Asp Lys Glu Leu Val Ser Asp Asp					
		1330		1335		1340	
35		Glu Glu Arg Gly Thr Gly Leu Glu Glu Asn Asn Gln Glu Glu Gln Ser					
		1345		1350		1355	1360
		Met Asp Ser Asn Leu Gly Glu Ala Ala Ser Gly Cys Glu Ser Glu Thr					
		1365		1370		1375	
		Ser Val Ser Glu Asp Cys Ser Gly Leu Ser Ser Gln Ser Asp Ile Leu					
		1380		1385		1390	
40		Thr Thr Gln Gln Arg Asp Thr Met Gln His Asn Leu Ile Lys Leu Gln					
		1395		1400		1405	
		Gln Glu Met Ala Glu Leu Glu Ala Val Leu Glu Gln His Gly Ser Gln					
		1410		1415		1420	
45		Pro Ser Asn Ser Tyr Pro Ser Ile Ile Ser Asp Ser Ser Ala Leu Glu					
		1425		1430		1435	1440
		Asp Leu Arg Asn Pro Glu Gln Ser Thr Ser Glu Lys Ala Val Leu Thr					
		1445		1450		1455	
		Ser Gln Lys Ser Ser Glu Tyr Pro Ile Ser Gln Asn Pro Glu Gly Leu					
		1460		1465		1470	
50		Ser Ala Asp Lys Phe Glu Val Ser Ala Asp Ser Ser Thr Ser Lys Asn					
		1475		1480		1485	
		Lys Glu Pro Gly Val Glu Arg Ser Ser Pro Ser Lys Cys Pro Ser Leu					
		1490		1495		1500	
55		Asp Asp Arg Trp Tyr Met His Ser Cys Ser Gly Ser Leu Gln Asn Arg					
		1505		1510		1515	1520
		Asn Tyr Pro Ser Gln Glu Glu Leu Ile Lys Val Val Asp Val Glu Glu					
		1525		1530		1535	
		Gln Gln Leu Glu Glu Ser Gly Pro His Asp Leu Thr Glu Thr Ser Tyr					
		1540		1545		1550	

Leu Pro Arg Gln Asp Leu Glu Gly Thr Pro Tyr Leu Glu Ser Gly Ile
 1555 1560 1565
 Ser Leu Phe Ser Asp Asp Pro Glu Ser Asp Pro Ser Glu Asp Arg Ala
 1570 1575 1580
 5 Pro Glu Ser Ala Arg Val Gly Asn Ile Pro Ser Ser Thr Ser Ala Leu
 1585 1590 1595 1600
 Lys Val Pro Gln Leu Lys Val Ala Glu Ser Ala Gln Ser Pro Ala Ala
 1605 1610 1615
 10 Ala His Thr Thr Asp Thr Ala Gly Tyr Asn Ala Met Glu Glu Ser Val
 1620 1625 1630
 Ser Arg Glu Lys Pro Glu Leu Thr Ala Ser Thr Glu Arg Val Asn Lys
 1635 1640 1645
 Arg Met Ser Met Val Val Ser Gly Leu Thr Pro Glu Glu Phe Met Leu
 1650 1655 1660
 15 Val Tyr Lys Phe Ala Arg Lys His His Ile Thr Leu Thr Asn Leu Ile
 1665 1670 1675 1680
 Thr Glu Glu Thr Thr His Val Val Met Lys Thr Asp Ala Glu Phe Val
 1685 1690 1695
 20 Cys Glu Arg Thr Leu Lys Tyr Phe Leu Gly Ile Ala Gly Gly Lys Trp
 1700 1705 1710
 Val Val Ser Tyr Phe Trp Val Thr Gln Ser Ile Lys Glu Arg Lys Met
 1715 1720 1725
 Leu Asn Glu His Asp Phe Glu Val Arg Gly Asp Val Val Asn Gly Arg
 1730 1735 1740
 25 Asn His Gln Gly Pro Lys Arg Ala Arg Glu Ser Gln Asp Arg Lys Ile
 1745 1750 1755 1760
 Phe Arg Gly Leu Glu Ile Cys Cys Tyr Gly Pro Phe Thr Asn Met Pro
 1765 1770 1775
 30 Thr Asp Gln Leu Glu Trp Met Val Gln Leu Cys Gly Ala Ser Val Val
 1780 1785 1790
 Lys Glu Leu Ser Ser Phe Thr Leu Gly Thr Gly Val His Pro Ile Val
 1795 1800 1805
 Val Val Gln Pro Asp Ala Trp Thr Glu Asp Asn Gly Phe His Ala Ile
 1810 1815 1820
 35 Gly Gln Met Cys Glu Ala Pro Val Val Thr Arg Glu Trp Val Leu Asp
 1825 1830 1835 1840
 Ser Val Ala Leu Tyr Gln Cys Gln Glu Leu Asp Thr Tyr Leu Ile Pro
 1845 1850 1855
 40 Gln Ile Pro His Ser His Tyr
 1860

45 <210> 2
 <211> 393
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> BINDING
 <222> (300)...(393)
 50 <223> /note="Binding site of BRCA1 protein"

 <300>
 <301> Harlow E., et al.
 <302> Molecular cloning and in vitro expression of a cDN
 55 <303> Mol. Cell. Biol.
 <304> 5
 <305> 7
 <306> 1601-1610
 <307> 1985-01-01

<308> Genbank K03199

<309> 1995-01-07

<400> 2

5 Met Glu Glu Pro Gln Ser Asp Pro Ser Val Glu Pro Pro Leu Ser Gln
 1 5 10 15
 Glu Thr Phe Ser Asp Leu Trp Lys Leu Leu Pro Glu Asn Asn Val Leu
 20 25 30
 Ser Pro Leu Pro Ser Gln Ala Met Asp Asp Leu Met Leu Ser Pro Asp
 10 35 40 45
 Asp Ile Glu Gln Trp Phe Thr Glu Asp Pro Gly Pro Asp Glu Ala Pro
 50 55 60
 Arg Met Pro Glu Ala Ala Pro Pro Val Ala Pro Ala Ala Pro
 65 70 75 80
 15 Thr Pro Ala Ala Pro Ala Pro Ala Pro Ser Trp Pro Leu Ser Ser Ser
 85 90 95
 Val Pro Ser Gln Lys Thr Tyr Gln Gly Ser Tyr Gly Phe Arg Leu Gly
 100 105 110
 Phe Leu His Ser Gly Thr Ala Lys Ser Val Thr Cys Thr Tyr Ser Pro
 115 120 125
 20 Ala Leu Asn Lys Met Phe Cys Gln Leu Ala Lys Thr Cys Pro Val Gln
 130 135 140
 Leu Trp Val Asp Ser Thr Pro Pro Pro Gly Thr Arg Val Arg Ala Met
 145 150 155 160
 25 Ala Ile Tyr Lys Gln Ser Gln His Met Thr Glu Val Val Arg Arg Cys
 165 170 175
 Pro His His Glu Arg Cys Ser Asp Ser Asp Gly Leu Ala Pro Pro Gln
 180 185 190
 His Leu Ile Arg Val Glu Gly Asn Leu Arg Val Glu Tyr Leu Asp Asp
 195 200 205
 30 Arg Asn Thr Phe Arg His Ser Val Val Val Pro Tyr Glu Pro Pro Glu
 210 215 220
 Val Gly Ser Asp Cys Thr Thr Ile His Tyr Asn Tyr Met Cys Asn Ser
 225 230 235 240
 35 Ser Cys Met Gly Gly Met Asn Arg Arg Pro Ile Leu Thr Ile Ile Thr
 245 250 255
 Leu Glu Asp Ser Ser Gly Asn Leu Leu Gly Arg Asn Ser Phe Glu Val
 260 265 270
 His Val Cys Ala Cys Pro Gly Arg Asp Arg Arg Thr Glu Glu Glu Asn
 275 280 285
 40 Leu Arg Lys Lys Gly Glu Pro His His Glu Leu Pro Pro Gly Ser Thr
 290 295 300
 Lys Arg Ala Leu Pro Asn Asn Thr Ser Ser Ser Pro Gln Pro Lys Lys
 305 310 315 320
 45 Lys Pro Leu Asp Gly Glu Tyr Phe Thr Leu Gln Ile Arg Gly Arg Glu
 325 330 335
 Arg Phe Glu Met Phe Arg Glu Leu Asn Glu Ala Leu Glu Leu Lys Asp
 340 345 350
 Ala Gln Ala Gly Lys Glu Pro Gly Gly Ser Arg Ala His Ser Ser His
 355 360 365
 50 Leu Lys Ser Lys Lys Gly Gln Ser Thr Ser Arg His Lys Lys Leu Met
 370 375 380
 Phe Lys Thr Glu Gly Pro Asp Ser Asp
 385 390

55

<210> 3

<211> 5143

<212> DNA

<213> Homo sapiens

<220>
 <221> TATA_signal
 <222> (4538)...(4541)

5 <221> protein_bind
 <222> (4441)...(4491)
 <223> /note="BRCA1 regulation site, corresponds to -93 -
 -143 of promoter."

10 <221> prim_transcript
 <222> (4584)...(44584)
 <223> /note="BRCA1 regulation site, corresponds to -93 -
 -143 of promoter."

15 <300>
 <301> El-Deiry W.S., et al.
 <302> Topological control of p21(WAF1/CIP1) expression i
 <303> Cancer Research
 <304> 55

20 <306> 2910-2919
 <307> 1995-07-01
 <308> Genbank U24170
 <309> 1995-07-20

25 <400> 3
 ggatccctgt agagatgctc aggctgctga ggagggcgcg gtgcttggtc tctatgaata 60
 cgatgaccta aagcaaaaaa agaagatggc tatgtcgggtg aagctctatg gaactgggga 120
 tcaggaggcc tggcagaaaag gagtctgtgt tgcttctggg cagaacttgg catgatggag 180
 acgccagcca gcgagatgat gccaacccaga tttgccgaaa ttattgagaa gaatctcaaa 240
 30 agccgctagt agtaaaccga gtttcatatc agaccaggt cttggattga ggaacaggca 300
 atgggatcat tcctcagtgt ggccaaagga tctgacgagc cctcagtctt cttggaaatt 360
 cactacatag gcagcccaa tgcagacaaa ccacccttg tttgttggga aaggaattac 420
 ctttgacagt ggtggtatct ccatcaaggc ttctgcaaat atggacctca tgagggccga 480
 catgggagga gctacaacta tatgtcagc cattgtgtct gctgcaaatc tcagtttgcc 540
 35 cattaatatt ataggtctgg ccctctgtg aaaacatgcc cagcggcaag gccacaagc 600
 tgggatgtt gttagagcca ggaacaggaa gaccatccag gttggtaca ctagtgta 660
 ggggaggctc atactggctg atgcgctctg ttactgtcac acatttaacc cgaaggtcat 720
 cctcaatgcc accaccttaa caggtgtcat agatgtagct ttggggtcag gtgccactgg 780
 ggtctttacc aattcatcct ggctctggaa caagctcttc gaggccagca ttgaaacagg 840
 40 ggaccgtgtc tggaggatgc ctctcttcaa acattgtaca agacagggtg tagattgcca 900
 gctggctgat gttaacaaca ttgaaaaata tagatctgcg ggagcatgta catctgcggc 960
 attcctgaaa gaattcgtga ctcacccata gtgggcacat ttagacatag caggtgtgat 1020
 gaccaacaaa gatgagggtc cctatctatg gaaaggcatg accgggaggc ccacaaggac 1080
 tctcatagag ttcttacttc gtttcagtca agacaatgct tagttcagat actcaaaaaa 1140
 45 tgtcttcaact ctgtcttaaa ttggacagtt gaagttaaaa ggtttttgaa tgaatggatg 1200
 aaaatatttt aaaggaggca atttatattt aaaaatgtag aacacaatga aatttttatg 1260
 ccttgatttt tttttcattt tacacaaaga tttatatatt ttttttttga gacggagtct 1320
 cactctgtca cccaggctgg agtgcagggtg gcatgatctc agctcactgc aacctccgcc 1380
 tcctagggtc aagcgattct cccacctcag ccacctgaat acctgggact acaggtgccc 1440
 50 accaccatgc cgggctgatt tttgtatttt taatggagac ggggtttcac catattggcc 1500
 aggctggtct caaaactcct gacctgtga tctgcccgcc tcggcctccc aaagtgtcgg- 1560
 gattacaggc gtaaacacc acccccggcc agtatatatt tttaattgag aagcaaaatt 1620
 gtacttcaga tttgtgatgc taggaacatg agcaaaactga aaattactaa ccacttgtca 1680
 gaaacaataa atccaacttt ttgtgcaaaa aaaaaaata caaatattag ctgggcatgg 1740
 55 tggtgcatgc ctgtaatccc agctactcgg gaggctgagg cagaattgct tgaacctggg 1800
 aggcggagac tgcagtgagc tgagattgtg ccactgctga ctttgtctca aaaaaaaaaa 1860
 caaaacaaaa aaacaaaatg aaaacaaaaa gccagggtct cctctgtctca ataattgtct 1920
 atctttgttc cgcctcttct ctgggggtct acttcttggg agcctgtgtg aaggtgaatt 1980
 cctctgaaag ctgactgccc ctatttggga ctccccagtc tctttctgag aaatggtgac 2040

	attgttccca	gcacttcctc	tcccttccta	ggcagcttct	gcagccacca	ctgagccttc	2100
	ctcacatcct	ccttcttcag	gcttgggctt	tccaccttcc	accattccccc	taccccatgc	2160
	tgctccaccg	cactctgggg	aggggggctg	actgggcact	cttgtccccc	aggctgagcc	2220
	tccctccatc	cctatgctgc	ctgcttccca	ggaacatgct	tgggcagcag	gctgtggctc	2280
5	tgattggctt	tctggccatc	aggaacatgt	cccaacatgt	tgagctctgg	catagaagag	2340
	gctgggtgct	atcttgcctt	tggtgtgcct	gttttcaggg	aggaaggagg	tggtaggaga	2400
	caggagacct	ctaaagaccc	caggtaaacc	ttagcctctt	actctgaaca	gggtatgtga	2460
	tctgccagca	ggatccttgc	gacagggtcg	ggatctgatg	catgtgtgct	tgtgtgagtg	2520
10	tggtctggga	gtcagattct	gtgtgtgact	tttaacagcc	tgctcccttg	ccttcttcag	2580
	ggcagaagtc	ctcccttaga	gtgtgtctgg	gtacacattc	aagtgcattg	ttgcaaaactt	2640
	ttttttttta	agcactgaat	agtactagac	acttagtagg	tacttaagaa	atattgaatg	2700
	tcgtgggtgt	ggtgagctag	aagttataaa	aaaaattctt	tcccaaaaac	aacaacaaaa	2760
	agaattattt	cattgtgaag	ctcagtacca	caaaaattca	ttacaataat	tcattacaag	2820
	cctttattaa	aaaaaatttt	ctcccaaaag	taaacagaca	gacaatgtct	agtctatttg	2880
15	aaatgcctga	aagcagaggg	gcttcaaggc	agtgggagaa	gggtgcctgtc	ctctgctgga	2940
	catttgacaa	ccagcccttt	ggatgggttg	tatgtatagg	agcgaagggtg	cagacagcag	3000
	tggggcttag	agtgggttcc	tgaggctgtg	ctgtggccct	tctgggggtt	agccacaatc	3060
	ctggcctgac	tccaggggcg	ggcaggccaa	gggggtctgc	tgctgtgtcc	tcccacccct	3120
	acctgggctc	ccatccccac	agcagaggag	aaagaagcct	gtcctccccg	aggtcagctg	3180
20	cgttagagta	agaagactgg	gcattgtctg	gcagagattt	ccagactctg	agcagcctga	3240
	gatgtcagta	attgtagctg	ctccaagcct	gggttctgtt	tttcagtggg	atttctgttc	3300
	agatgaacaa	tccatcctct	gcaatttttt	aaaagcaaaa	ctgcaaatgt	ttcaggcaca	3360
	gaaaggaggg	aaagggtgaag	tccaggggag	gtcaggggtg	tgaggtagat	gggagcggat	3420
	agacacatca	ctcatttctg	tgtctgtcag	aagaaccagt	agacacttcc	agaattgtcc	3480
25	tttattttat	tcattctccat	aaaccatctg	caaatgaggg	ttatttggca	tttttgtcat	3540
	tttggaaacca	cagaaataaa	ggatgacaag	cagagagccc	cgggcaggag	gcaaaagtcc	3600
	tgtgttccaa	ctatagtcac	ttctttgctg	catgatctga	gttaggtcac	cagacttctc	3660
	tgagccccag	tttccccagc	agtgtatagc	ggctatgtgg	ggagtattca	ggagacagac	3720
	aactcactcg	tcaaactctc	cccttcctgg	ccaacaaagc	tgctgcaacc	acagggtttt	3780
30	cttctgttca	ggtgagtgtg	gggtgtaggg	agattgggtc	aatgtccaat	tcttctgttt	3840
	ccctggagat	caggttgccc	ttttttggta	gtctctccaa	ttccctcctt	cccgggaagca	3900
	tgtgacaatc	aacaactttg	tatacttaag	ttcagtggac	ctcaatttcc	tcattctgtg	3960
	aataaacggg	actgaaaaat	cattctggcc	tcaagatgct	ttgttggggg	gtctaggtgc	4020
	tccaggtgct	tctgggagag	gtgacctagt	gagggatcag	tggaataga	ggtgatattg	4080
35	tggggctttt	ctggaaattg	cagagaggtg	catcgttttt	ataatttatg	aatttttatg	4140
	tattaatgtc	atcctcctga	tcttttcagc	tgcatggggg	aaatccttgc	ctgccagagt	4200
	gggtcagcgg	tgagccagaa	agggggctca	ttctaacagt	gctgtgtcct	cctggagagt	4260
	gccaactcat	tctccaagta	aaaaaagcca	gatttgtggc	tcacttcgtg	gggaaatgtg	4320
	tccagcgcac	caacgcaggc	gagggactgg	gggaggaggg	aagtgccctc	ctgcagcacg	4380
40	cgagggttcc	ggaccggctg	gcctgctgga	actcggccag	gctcagctgc	tccgcgctgg	4440
	gcagccagga	gcctggggcc	cggggagggc	ggtcccgggc	ggcgcgggtg	gccgagcgcg	4500
	ggtcgcctcc	ttgaggcggg	cccgggcggg	gcggttgtat	atcagggccg	cgctgagctg	4560
	cgccagctga	ggtgtgagca	gctgccgaag	tcagttcctt	gtggagccgg	agctgggcgc	4620
	ggattcgccg	aggcaccgag	gcactcagag	gaggtgagag	agcggcggca	gacaacaggg	4680
45	gaccccgggc	cggcgggcca	gagccgagcc	aagcgtgccc	gcgtgtgtcc	ctgcgtgtcc	4740
	gcgaggatgc	gtgttcgcgg	gtgtgtgctg	cgttcacagg	tgtttctgcg	gcaggtgaat	4800
	gacgggcgtg	ggtcggtgcg	cgctcggtct	gcgcacacgg	tgtctctaag	tgccggtgtg	4860
	acgagggctg	ggatgtgccc	gagaccccg	gcggagagcg	ggattacaag	tacaggaatc	4920
	cctggccacg	ctccccgcc	ctggaaaccc	agctggggcg	agggagggcg	tggacgggac	4980
50	cgttctggga	gctcgctttt	ggctgcggtt	ggctccaggc	cccaggcgca	gtttgctcgc	5040
	ggcgtgggga	tgaagtcctg	gtccctggag	gggcccagga	agggcgagga	aagcggagtg	5100
	gagtaagttc	gtctaggatc	ggctccgggt	ggctctggga	tcc		5143

55

<210> 4

<211> 35

<212> DNA

<213> Artificial Sequence

<220>
<223> Sequence source:/note="Synthetic construct"

<400> 4
5 gcaagcttgc caccatggat ttatctgctc ttcgc 35

<210> 5
<211> 21
<212> DNA
10 <213> Artificial Sequence

<220>
<223> Sequence source:/note="Synthetic construct"

<400> 5
15 ttgtgagggg acgctcttgt a 21

<210> 6
<211> 21
20 <212> DNA
<213> Artificial Sequence

<220>
<223> Sequence source:/note="Synthetic construct"

<400> 6
25 aggagatgtg gtcaatggaa g 21

<210> 7
30 <211> 50
<212> DNA
<213> Artificial Sequence

<220>
35 <223> Sequence source:/note="Synthetic construct"

<400> 7
40 tatcatgatg taggtctcct tttacgcttt aatttattgt agtggctgtg 50

<210> 8
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
45 <223> Sequence source:/note="Synthetic construct"

<400> 8
50 aagcttgcca ccatggagga gccgcagtca 30

<210> 9
55 <211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Sequence source:/note="Synthetic construct"

<400> 9

atgcggccgc tcagtctgag tcag 24

5 <210> 10
<211> 31
<212> DNA
<213> Artificial Sequence

10 <220>
<223> Sequence source:/note="Synthetic construct"

<400> 10
taagcttgcc accatggctg cttgtgaatt t 31

15 <210> 11
<211> 25
<212> DNA
<213> Artificial Sequence

20 <220>
<223> Sequence source:/note="Synthetic construct"

<400> 11
actcgagtca taagtttgaa tccat 25

25 <210> 12
<211> 35
<212> DNA
<213> Artificial Sequence

30 <220>
<223> Sequence source:/note="Synthetic construct"

<400> 12
gcaagcttgc caccatggat ttatctgctc ttcgc 35

35 <210> 13
<211> 20
<212> DNA
<213> Artificial Sequence

40 <220>
<223> Sequence source:/note="Synthetic construct"

<400> 13
45 ggagcccact tcattagtag 20

<210> 14
<211> 35
<212> DNA
50 <213> Artificial Sequence

<220>
<223> Sequence source:/note="Synthetic construct"

55 <400> 14
gcaagcttgc caccatggat ttatctgctc ttcgc 35

<210> 15
<211> 23

<212> DNA
<213> Artificial Sequence

<220>
5 <223> Sequence source:/note="Synthetic construct"

<400> 15
acctgtgtca agctgaaaag cac 23

10 <210> 16
<211> 30
<212> DNA
<213> Artificial Sequence

15 <220>
<223> Sequence source:/note="Synthetic construct"

<400> 16
20 aagcttgcca ccatgtgtgc ttttcagctt 30

<210> 17
<211> 25
<212> DNA
<213> Artificial Sequence

25 <220>
<223> Sequence source:/note="Synthetic construct"

<400> 17
30 tctcgagtca atttgtgagg ggacg 25

<210> 18
<211> 31
<212> DNA
35 <213> Artificial Sequence

<220>
<223> Sequence source:/note="Synthetic construct"

<400> 18
40 taagcttgcc accatggctg cttgtgaatt t 31

<210> 19
<211> 25
45 <212> DNA
<213> Artificial Sequence

<220>
<223> Sequence source:/note="Synthetic construct"

50 <400> 19
tctcgagtca atttgtgagg ggacg 25

<210> 20
55 <211> 27
<212> DNA
<213> Artificial Sequence

<220>

<223> Sequence source:/note="Synthetic construct"

<400> 20
5 gtcgaattcg aggagccgca gtcagat 27

<210> 21
<211> 24
<212> DNA
10 <213> Artificial Sequence

<220>
<223> Sequence source:/note="Synthetic construct"

<400> 21
15 atgcgggccgc tcagtctgag tcag 24

<210> 22
<211> 27
<212> DNA
20 <213> Artificial Sequence

<220>
<223> Sequence source:/note="Synthetic construct"

<400> 22
25 gtcgaattcg aggagccgca gtcagat 27

<210> 23
<211> 25
30 <212> DNA
<213> Artificial Sequence

<220>
<223> Sequence source:/note="Synthetic construct"

<400> 23
35 atgcgggccgc caaatcatcc attgc 25

<210> 24
40 <211> 27
<212> DNA
<213> Artificial Sequence

<220>
45 <223> Sequence source:/note="Synthetic construct"

<400> 24
50 gtcgaattcg aggagccgca gtcagat 27

<210> 25
<211> 26
<212> DNA
55 <213> Artificial Sequence

<220>
<223> Sequence source:/note="Synthetic construct"

<400> 25
tactcgagat ttccttcac tcggat 26

	<210> 26	
	<211> 23	
	<212> DNA	
5	<213> Artificial Sequence	
	<220>	
	<223> Sequence source:/note="Synthetic construct"	
	<400> 26	
10	atgaattcca gaaaacctac cag	23
	<210> 27	
	<211> 22	
	<212> DNA	
15	<213> Artificial Sequence	
	<220>	
	<223> Sequence source:/note="Synthetic construct"	
	<400> 27	
20	tactcgaggg cagctcgtgg tg	22
	<210> 28	
	<211> 26	
25	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Sequence source:/note="Synthetic construct"	
30	<400> 28	
	atgaattctt gcgtgtggag tatattg	26
	<210> 29	
35	<211> 24	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
40	<223> Sequence source:/note="Synthetic construct"	
	<400> 29	
	atgcggccgc tcagtctgag tcag	24
45	<210> 30	
	<211> 23	
	<212> DNA	
	<213> Artificial Sequence	
50	<220>	
	<223> Sequence source:/note="Synthetic construct"	
	<400> 30	
55	atgaattccc cccagggagc act	23
	<210> 31	
	<211> 23	
	<212> DNA	
	<213> Artificial Sequence	

<220>
<223> Sequence source:/note="Synthetic construct"

<400> 31
5 tgcgggccgct cagtctgagt cag 23

<210> 32
<211> 22
<212> DNA
10 <213> Artificial Sequence

<220>
<223> Sequence source:/note="Synthetic construct"

<400> 32
15 agaattctgt gcttttcagc tt 22

<210> 33
<211> 25
20 <212> DNA
<213> Artificial Sequence

<220>
<223> Sequence source:/note="Synthetic construct"

<400> 33
25 tctcgagtca atttgtgagg ggacg 25

30

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/07150

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07K 1/00; C07H 21/02

US CL : 530/350; 536/23.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/350; 536/23.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Genbank, SEQ ID NO:1 and 3, p21 gene, isolated, promoter, upstream p53

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GENBANK Sequence Database, National Center for Biotechnology Information, National Library of Medicine, (Bethesda, Maryland, USA) Accession U24170, El-Deiry et al., 'Topological Control of P21 WAF1/CIP1 Expression in Normal and Neoplastic Tissues,' abstract, Cancer Res. 55 (13), 2910-2919, 20 July 1995, see entire sequence listing.	1-6
X	EL-DEIRY et al. Topological Control of P21 WAF1/CIP1 Expression in Normal and Neoplastic Tissues. Cancer Research. July 1995, Vol. 55, pages 2910-2919, see entire article.	1-6
X	MIKI et al. A Strong Candidate for the Breast and Ovarian Cancer Susceptibility Gene BRCA1. Science. 07 October 1994, Vol. 266, pages 66-71, see especially Figure 2, page 67.	7-9

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

21 JUNE 1999

Date of mailing of the international search report

28 JUL 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

SUSAN UNGAR

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/07150**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
1-9
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/07150

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-6, drawn to an isolated nucleic acid molecule comprising a region of an upstream promotor of a p21 gene which contains an element required for p53-independent BRCA1-mediated transactivation of the gene.
Group II, claim(s) 7-9, drawn to an isolated polypeptide comprising a region of BRCA1 protein.
Group III, claim(s) 10, drawn to an antibody specific for BRCA1 protein.
Group IV, claim(s) 11-12, drawn to an isolated nucleic acid encoding BRCA1 protein and a vector comprising said nucleic acid.
Group V, claim(s) 13-14, drawn to an isolated polypeptide comprising a region of p53.
Group VI, claim(s) 16, drawn to an antibody specific for p53.
Group VII, claim(s) 17-18, drawn to a nucleic acid molecule encoding p53 and a vector comprising said nucleic acid.
Group VIII, claim(s) 19-20, drawn to an isolated protein-protein complex comprising p53 and BRCA1.
Group IX, claim(s) 21-23 and 36, drawn to a method of determining if a test compound modulates the formation of a complex comprising BRCA1 and p53.
Group X, claim(s) 24, 25, 37 and 38, drawn to a method for determining if a test compound enhances or interferes with p53-independent BRCA1-mediated transactivation of a p21 gene.
Group XI, claim(s) 26-28 and 39-40, drawn to a method for determining if a test compound modulates tumor suppression resulting from p53-independent BRCA1-mediated transactivation of a p21 gene.
Group XII, claim(s) 29-30 and 41-42, drawn to a method for determining if a test compound modulates p53-independent BRCA1-mediated transactivation of a gene.
Group XIII, claim(s) 31-35 and 43-44, drawn to a method for determining if a test compound modulates tumor suppression resulting from p53-dependent BRCA1-mediated transactivation of a tumor suppressor gene.
Group XIV, claim(s) 45-46, drawn to a method for diagnosing predisposition for developing a tumor.
Group XV, claim(s) 47, drawn to a method for enhancing tumor suppression.
Group XVI, claim 48, drawn to a method for increasing cell proliferation.

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack Unity of Invention because they are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for more than one species to be searched, the appropriate additional search fees must be paid. The species are as follows:

Group XIV

Species A, Claims 45 and 46, a method for diagnosing predisposition for developing a tumor comprising detecting defects in the subjects BRCA1 genes.

Species B, Claims 45 and 46, a method for diagnosing predisposition for developing a tumor comprising detecting defects in the subjects BRCA1 proteins.

Species C, Claims 45 and 46, a method for diagnosing predisposition for developing a tumor comprising detecting defects in the subjects p21 promoters.

Species D, Claim 46, a method for diagnosing predisposition for developing a tumor comprising detecting defects in the subjects p53 genes.

Species E, Claim 46, a method for diagnosing predisposition for developing a tumor comprising detecting defects in the subjects p53 proteins.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/07150

The inventions listed as Groups I-XVI do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The technical feature linking groups I-XVI appears to be an isolated nucleic acid molecule comprising a region of an upstream promoter of a p21 gene that contains an element required for p53-independent BRCA1-mediated transactivation of the gene, wherein the region comprises nucleotides 4441-4591 of SEQ ID NO:3.

However, El-Deiry et al(HSU24170), Genbank Sequence Database (Accession U24170), National Center for Biotechnology Information, National Library of Medicine, Bethesda, Maryland, publicly available 20 July 1995 teaches an isolated nucleic acid molecule which has 100% identity to SEQ ID NO:3.

Therefore, the technical feature linking Groups I-XVI is not considered to be a special technical feature as defined by PCT Rule 13.2 as it does not define a contribution over the prior art.

The special technical feature of Group I is considered to be an isolated nucleic acid molecule comprising a p53-independent p21 promoter.

The special technical feature of Group II is considered to be a BRAC1 protein.

The special technical feature of Group III is considered to be an antibody specific for BRAC1.

The special technical feature of Group IV is considered to be a nucleic acid encoding BRAC1.

The special technical feature of Group V is considered to be a p53 protein.

The special technical feature of Group VI is considered to be an antibody specific for p53.

The special technical feature of Group VII is considered to be a nucleic acid encoding p53.

The special technical feature of Group VIII is considered to be a protein complex.

The special technical feature of Group IX is considered to be a method of determining if a test compound modulates formation of a BRCA1/p53 complex.

The special technical feature of Group X is considered to be a method for determining if a test compound modulates p53-independent BRCA1-mediated transactivation of a p21 gene.

The special technical feature of Group XI is considered to be a method for determining if a test compound modulates p53-independent BRCA-1 mediated transactivation of a p21 gene.

The special technical feature of Group XII is considered to be a method for determining if a test compound modulates p53-independent BRCA-1 mediated transactivation of a p21 gene.

The special technical feature of Group XIII is considered to be a method for determining if a test compound modulates tumor suppression.

The special technical feature of Group XIV is considered to be a method for diagnosis of predisposition for developing a tumor.

The special technical feature of Group XV is considered to be a method for enhancing tumor suppression.

The special technical feature of Group XVI is considered to be a method for increasing cell proliferation.

Accordingly, Groups I-XVI are not so linked by the same or a corresponding special technical feature as to form a single general inventive concept.

The species listed above do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons:

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/07150

The species as claimed are materially distinct methods which differ at least in objectives, method steps, reagents and/or dosages and/or schedules used, response variables, and criteria for success.